

# $\alpha$ -Pinene, a Major Constituent of Pine Tree Oils, Enhances Non-Rapid Eye Movement Sleep in Mice through GABA<sub>A</sub>-benzodiazepine Receptors<sup>[S]</sup>

Hyejin Yang, Junsung Woo, Ae Nim Pae, Min Young Um, Nam-Chul Cho, Ki Duk Park, Minseok Yoon, Jiyoung Kim, C. Justin Lee, and Suengmok Cho

*Division of Functional Food Research, Korea Food Research Institute, Seongnam, Republic of Korea (H.Y., M.Y.U., M.Y., J.K., S.C.); Center for Neuroscience and Functional Connectomics, Korea Institute of Science and Technology, Seoul, Republic of Korea (J.W., C.J.L.); Convergence Research Center for Diagnosis, Treatment and Care System of Dementia, Seoul, Republic of Korea (A.N.P., N.-C.C., K.D.P.); and KU-KIST Graduate School of Converging Sciences and Technologies, Korea University, Seoul, Republic of Korea (C.J.L.)*

Received May 9, 2016; accepted August 19, 2016

## ABSTRACT

$\alpha$ -Pinene is a major monoterpene of the pine tree essential oils. It has been reported that  $\alpha$ -pinene shows anxiolytic and hypnotic effects upon inhaled administration. However, hypnotic effect by oral supplementation and the molecular mechanism of  $\alpha$ -pinene have not been determined yet. By combining *in vivo* sleep behavior, *ex vivo* electrophysiological recording from brain slices, and *in silico* molecular modeling, we demonstrate that ( $-$ )- $\alpha$ -pinene shows sleep enhancing property through a direct binding to GABA<sub>A</sub>-benzodiazepine (BZD) receptors by acting as a partial modulator at the BZD binding site. The effect of ( $-$ )- $\alpha$ -pinene on sleep-wake profiles was evaluated by recording electroencephalogram and electromyogram. The molecular mechanism of ( $-$ )- $\alpha$ -pinene was investigated by electrophysiology and molecular docking study. ( $-$ )- $\alpha$ -pinene significantly increased the duration of non-rapid eye movement sleep (NREMS) and

reduced the sleep latency by oral administration without affecting duration of rapid eye movement sleep and delta activity. ( $-$ )- $\alpha$ -pinene potentiated the GABA<sub>A</sub> receptor-mediated synaptic response by increasing the decay time constant of sIPSCs in hippocampal CA1 pyramidal neurons. These effects of ( $-$ )- $\alpha$ -pinene on sleep and inhibitory synaptic response were mimicked by zolpidem, acting as a modulator for GABA<sub>A</sub>-BZD receptors, and fully antagonized by flumazenil, an antagonist for GABA<sub>A</sub>-BZD receptor. ( $-$ )- $\alpha$ -pinene was found to bind to aromatic residues of  $\alpha$ 1- and  $\gamma$ 2 subunits of GABA<sub>A</sub>-BZD receptors in the molecular model. We conclude that ( $-$ )- $\alpha$ -pinene enhances the quantity of NREMS without affecting the intensity of NREMS by prolonging GABAergic synaptic transmission, acting as a partial modulator of GABA<sub>A</sub>-BZD receptors and directly binding to the BZD binding site of GABA<sub>A</sub> receptor.

## Introduction

Pine trees of the genus *Pinus* comprise of more than 100–250 species and are widely spread out all over the world (Judzentiene and Kupcinskiene, 2008; Inoannou et al., 2014). It has been one of the most popular plants worldwide because of their medicinal and aromatic properties (Lee et al., 2005; Yang et al., 2010). It has been accepted that these traditional and pharmaceutical uses of pines are due to their essential oils (Hmamouchi et al., 2001). Pine essential oils possess numerous

biologic activities, such as anti-inflammatory, antimicrobial, analgesic, and antistress effects (Süntar et al., 2012; Xie et al., 2015). They are mainly composed of monoterpene such as  $\alpha$ - and  $\beta$ -pinene, 3-carene, limonene, and terpinene (Judzentiene and Kupcinskiene, 2008).

$\alpha$ -Pinene [2,6,6-trimethylbicyclo(3.1.1)-2-hept-2-ene] is the major monoterpene of pine essential oils (Groot and MacDonald, 2002) and a hydrocarbon group of bicyclic terpenes with a strong turpentine odor (Bakkali et al., 2008). It has been widely used as a food flavoring ingredient (Limberger et al., 2007; Rivas et al., 2012) and was approved as a food additive generally recognized as safe by U.S. Food and Drug Administration (FDA, 2015). In addition, a number of studies have attributed biological properties, including antimicrobial (Gomes-Carneiro et al., 2005), hypertensive (Kamal et al., 2003), antinociceptive (Him et al., 2008), and anti-inflammatory (Orhan et al., 2006) effects to  $\alpha$ -pinene.

Recently, Satou et al. (2014) reported that inhalation of  $\alpha$ -pinene produces anxiolytic activity in an elevated plus maze

This study was supported by grants from the Korea Food Research Institute [Grant E0164503-01], Creative Research Initiative Program, Korean National Research Foundation [Grant 2015R1A3A2066619], Korea Institute of Science and Technology [Grant 2E26662], KU-KIST Graduate School of Science and Technology program [Grant R1435281], and National Research Council of Science and Technology (NST) grant by the Korea government (MSIP) [Grant CRC-15-04-KIST].

H.Y. and J.W. contributed equally to this work.  
[dx.doi.org/10.1124/mol.116.105080](http://dx.doi.org/10.1124/mol.116.105080).

[S] This article has supplemental material available at [molpharm.aspetjournals.org](http://molpharm.aspetjournals.org).

**ABBREVIATIONS:** ACSF, artificial cerebrospinal fluid; BZD, benzodiazepine; EEG, electroencephalogram; EMG, electromyogram; ICR, imprinting control region; NREMS, non-REM sleep; REMS, rapid eye movement sleep; sIPSCs, spontaneous inhibitory postsynaptic currents; Wake, wakefulness.

test in mice. They also confirmed the accumulation of  $\alpha$ -pinene in the brain. According to Yamaoka et al. (2005), inhalation of  $\alpha$ -pinene significantly increased rapid eye movement sleep in rats. Despite the popular usage of  $\alpha$ -pinene as a food ingredient and therapeutic agent, its action on sleep and anxiety, the related molecular mechanism, and effect by oral supplementation of  $\alpha$ -pinene have not been determined yet. In this study, we investigated the hypnotic effect of orally administrated  $\alpha$ -pinene. Using pharmacological tools (e.g., zolpidem and flumazenil), electrophysiology, and molecular modeling, we set out to identify the molecular mechanism of  $\alpha$ -pinene.

## Materials and Methods

### Materials

(-)- $\alpha$ -Pinene (CAS no. 7785-26-4) was purchased from Santa Cruz Biotechnology Inc. (Dallas, TX). Zolpidem (CAS no. 82626-48-0), a GABA<sub>A</sub>-benzodiazepine (BZD) receptor agonist, was used as a reference hypnotic drug and obtained from Ministry of Food and Drug Safety of Korea (Cheongwon-gun, Chungcheongbuk-do, Korea). Flumazenil (CAS no. 78755-81-4), an antagonist of GABA<sub>A</sub>-BZD receptors, was purchased from Sigma-Aldrich Inc. (St. Louis, MO). Molecular structures and weights of (-)- $\alpha$ -pinene, zolpidem, and flumazenil are shown in Fig. 1. All other chemicals and reagents were of the highest grade available.

### Animals

All procedures involving animals were conducted in accordance with the animal care and use guidelines of the Korea Food Research Institutional Animal Care and Use Committee (permission number: KFRI-M-12027). Imprinting control region (ICR; male, 18–22 g) and C57BL/6N (male 27–30 g) mice were purchased from Koatech Animal Inc. (Pyeongtaek, Korea). The animals were housed in an insulated, sound-proof recording room maintained at an ambient temperature of  $23 \pm 0.5^\circ\text{C}$ , with a constant relative humidity ( $55 \pm 2\%$ ) on an automatically controlled 12-hour light/12-hour dark cycle (lights off at 1700). They had free access to food and water. All efforts were made to minimize animal suffering and to use only the number of animals required for the production of reliable scientific data.

### Pentobarbital-Induced Sleep Test


The initial screening for hypnotic effect of (-)- $\alpha$ -pinene sleep was done with pentobarbital-induced sleep, as previously described (Cho et al., 2011). Experiment was performed between 1300 and 1700, and the ICR mice were fasted for 24 hours before the experiment to minimize the drowsiness induced by food. (-)- $\alpha$ -Pinene and zolpidem were administered orally to the ICR mice ( $n = 10$ ) 45 minutes before the pentobarbital injection (45 mg/kg, i.p.). After the injection (intraperitoneal) of pentobarbital, mice were placed in individual cages and observed for measurements of sleep latency

and duration. The observers were blinded to the individual treatments. The mice were considered asleep if they stayed immobile and lost their righting reflexes when positioned on the back. The sleep latency was defined as the elapsed time from pentobarbital injection to onset of righting reflex loss. The sleep duration was defined as the difference in time between the loss and the recovery of the righting reflex.

### Analysis of Sleep Architecture

**Pharmacological Treatments.** (-)- $\alpha$ -Pinene was dissolved in sterile saline containing 5% Tween 80 immediately before use and administered orally to the C57BL/6N mice (each group,  $n = 8$ ) at 1700 on the experimental day at a dose of 25, 50, or 100 mg/kg. The positive control zolpidem (10 mg/kg) was administered in the same manner as (-)- $\alpha$ -pinene. Flumazenil was dissolved in sterile saline and injected intraperitoneally 15 minutes before (-)- $\alpha$ -pinene or zolpidem administration. For baseline data, mice were injected with the vehicle (saline containing 5% Tween 80) at 1645 (intraperitoneally) and 1700 (orally).

**Polygraphic Recordings and Vigilance State Analysis.** Under pentobarbital anesthesia (50 mg/kg, i.p.), the C57BL/6N mice were chronically implanted with a head mount (#8201, Pinnacle Technology Inc., Lawrence, KS) installed with electroencephalogram (EEG) and electromyogram (EMG) electrodes for polysomnographic recordings. The front edge of the head mount was placed 3.0 mm anterior to bregma, and four electrode screws for EEG recording were positioned in holes perforated into the skull. Two EMG wire electrodes were inserted into the nuchal muscles. The head mount was fixed to the skull with dental cement. After surgery, mice were allowed to recover in individual cages for 1 week and habituated to the recording conditions for 3–4 days before the experiment. The EEG and EMG recordings were carried out by means of a slip ring designed so that the movement of the mice was not restricted. EEG and EMG were recorded using the PAL-8200 data acquisition system (Pinnacle Technology Inc.). The EEG and EMG signals were amplified (100 $\times$ ), filtered (low-pass filter: 25 Hz EEG and 100 Hz EMG), and stored at a sampling rate of 200 Hz. Sleep states were monitored for a period of 48 hours, which comprised baseline and experimental days. Baseline recordings were taken for each animal during 24 hours, beginning at 1700. These baseline recordings served as controls for the same animal. The mice were considered asleep showing no EMG signal. The vigilance states were automatically classified by a 10-second epoch as wakefulness (Wake), rapid eye movement sleep (REMS), or non-REM sleep (NREMS) by SleepSign ver. 3.0 (Kissei Comtec, Nagano, Japan). As a final step, defined sleep-wake stages were examined visually and corrected if necessary. The sleep latency was defined as the time from drug administration to the appearance of the first NREMS episode lasting for at least 20 seconds. Bouts of NREMS, REMS, and Wake were defined as periods of one or more consecutive epochs (each epoch: 10 seconds). Each delta power of NREMS in the range of 0.5–4 Hz was first summated and then normalized as a percentage of the corresponding mean delta power of NREMS.



**Fig. 1.** Chemical structures and molecular weight (MW) of (-)- $\alpha$ -pinene, zolpidem, and flumazenil.

## Electrophysiological Measurement

**Slice Preparation.** Adult mice (7–9 weeks) were deeply anesthetized until cessation of breathing and subsequently decapitated. The brain was rapidly removed and submerged in an ice-cold oxygenated artificial cerebrospinal fluid (ACSF) composed of (in mM) 130 NaCl, 24 NaHCO<sub>3</sub>, 3.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 1 CaCl<sub>2</sub>, 3 MgCl<sub>2</sub>, 10 glucose at pH 7.4, and was bubbled with 5% CO<sub>2</sub>/95% O<sub>2</sub>. Transverse mouse brain slices (300 μm) containing hippocampus were acutely prepared with a vibratome (DSK LinearSlicer, Kyoto, Japan), and incubated in a chamber with oxygenated ACSF at room temperature for 1 hour before use.

**Recording of sIPSCs.** The standard ACSF recording solution was composed of (mM): 130 NaCl, 24 NaHCO<sub>3</sub>, 3.5 KCl, 1.25 NaH<sub>2</sub>PO<sub>4</sub>, 1.5 CaCl<sub>2</sub>, 1.5 MgCl<sub>2</sub>, and 10 glucose saturated with 95% O<sub>2</sub>-5% CO<sub>2</sub> at pH 7.4. The internal solution was composed of (mM): 140 CsCl, 10 EGTA, 10 HEPES, 4 Mg-ATP, 10 QX-314. To block the spontaneous EPSC, APV (2-amino-5-phosphonopentanoic acid; 50 μM; Tocris, Minneapolis, MN) and CNQX (6-cyano-7-nitroquinoxaline-2,3-dione; 20 μM; Tocris) were added into ACSF. Recordings were obtained using Axopatch 200A (Axon Instruments, Union City, CA) and filtered at 2 kHz. In case of sIPSC recording, recordings were digitized at 10 kHz, and analyzed using pCLAMP 10 (Molecular Devices, Sunnyvale, CA) and Mini Analysis Program (Synaptosoft, Decatur, GA). The sIPSCs were automatically detected. All experimental procedures described were performed in accordance with the institutional guidelines of Korea Institute of Science and Technology (KIST, Seoul, Korea).

## Molecular Modeling


**Homology Modeling.** Homology modeling was performed with MODELER in DiscoveryStudio program (Accelrys, San Diego, CA). The X-ray crystal structure of a human GABAA receptor, the β3 homopentamer (PDB code: 4COF) was employed as template for the most abundant α1β2γ2 subtype of GABAA receptor (Miller and Aricescu, 2014). The sequences of the human α1 (P14867), β2 (P47870), and γ2 (P18507) were retrieved from the UniProt database (<http://www.uniprot.org>) and aligned with ClustalW implemented in DiscoveryStudio program. The mismatch residues between α1- and β3-subunits were manually edited to remove gaps following the sequence alignment reported by Miller and Aricescu (2014). The sequence identities of α1, β2, and γ2 with β3 were 40.6, 89.8, and 41.4%, and the sequence similarities of them were 62.6, 95.8, and 63.7%. Procheck validation indicated that 1669 of total 1670 residues of homology model was in allowed region of Ramachandran plot, and only one residue Arg204 of α1-subunit was an outlier. The cocrystallized ligands containing benzamidine, sugars, and ions were removed and the best model of α1β2γ2 subtype GABA<sub>A</sub> receptor among 50 generated models having various conformations with MODELER was selected by optimally satisfying spatial restraints derived from the alignment and expressed as probability density functions for the features restrained, which is calculated from the relationship of Cα-Cα pairs and main-chain dihedral angles between homology model and template as spatial restraints (Sali and Blundell, 1993).

**Molecular Docking Study.** Molecular Docking was performed using Schrodinger package program (Schrodinger LLC, New York, NY). Homology model of α1β2γ2 subtype GABAA receptor was neutralized and energetically minimized with OPLS2005 forcefield using Protein Prep Wizard. Parameters of Protein Prep Wizard were set to default value. Chemical structures of (-)-α-pinene, zolpidem, and flumazenil were sketched using ChemDraw program. Ligands were prepared with protonation at pH 7.4 and energy minimization with OPLS2005 forcefield using LigPrep module. Parameters of LigPrep were set to default value. Binding modes of ligands were predicted into benzodiazepine (BZD) site of GABAA receptor using InduceFit docking module. The grid box was automatically set into the centroid region of α1Y209, α1H101, α1Y159, γ2F77, γ2M130, and

γ2T142 residues. Glide XP docking algorithm was used for more extensive thorough torsional refinement and sampling. Other parameters were default. Predicted binding poses of compound were selected with low Gscore for interactions. Gscore is an empirical scoring function that includes interaction energies of hydrogen bond, hydrophobic, van-der Waals in binding site, and ligand strain energy.

## Data Analysis



All data were expressed as the mean ± S.E.M. (standard error of mean). Statistical analysis was performed with the Prism 5.0 (GraphPad Software Inc., San Diego, CA). For multiple comparisons, data were analyzed using one-way analysis of variance followed by



**Fig. 2.** Experimental procedure for pentobarbital-induced sleep test (A). Effects of the administration of (-)-α-pinene alone and with flumazenil on sleep latency (B) and sleep duration (C) in pentobarbital-treated (45 mg/kg, i.p.) ICR mice. The CON (5% Tween 80-saline, 10 ml/kg), zolpidem, and (-)-α-pinene were administered orally to mice 45 minutes before injection (intraperitoneal) of pentobarbital. Flumazenil was intraperitoneally injected (intraperitoneally) 15 minutes before oral administration of CON, zolpidem, and (-)-α-pinene. Each value, the mean ± S.E.M. was calculated from 10 mice. \**P* < 0.05, \*\**P* < 0.01, significant compared with the CON (Dunnett's test). ##*P* < 0.01, significant between the flumazenil treatment and nonflumazenil treatment (unpaired Student's *t* test). CON, control group; NS, not significant.

## A Experimental procedure for polygraphic recordings

Animal (n=8): C57BL/6N mice



**Fig. 3.** Experimental procedure for analysis of sleep architecture in C57BL/6N mouse (A). Typical electroencephalogram (EEG) and electromyogram (EMG) waveforms (B) and fast Fourier transform (FFT) spectrum (C) in states of Wake, REMS, and NREMS.

Dunnett's test. Comparisons between two-group data were analyzed by the unpaired Student's *t* test. The significance level was set at  $P < 0.05$  for all statistical tests.

## Results

### Effects of (–)- $\alpha$ -Pinene in the Pentobarbital-Induced Sleep Test in ICR Mice


To investigate whether oral administration of (–)- $\alpha$ -pinene (Fig. 1) produces sedative-hypnotic effects, we first used the pentobarbital-induced sleep test in ICR mice. As expected, a well-known hypnotic drug, zolpidem (Fig. 1; 10 mg/kg, oral) significantly ( $P < 0.01$ ) potentiated pentobarbital-induced sleep in mice relative to the control group (Fig. 2). (–)- $\alpha$ -Pinene (12.5, 25, 50, and 100 mg/kg, p.o.) also decreased sleep latency and increased sleep duration in a dose-dependent manner. In particular, administration of 100 mg/kg of (–)- $\alpha$ -pinene was found to prolong sleep duration up to  $114.7 \pm 8.2$  minutes to the level similar to that of zolpidem at 10 mg/kg ( $124.2 \pm 4.9$  minutes).

### Effects of (–)- $\alpha$ -Pinene on Sleep Architecture in C57BL/6N Mice: Sleep Latency and Amounts of NREMS and REMS

To determine the effects of (–)- $\alpha$ -pinene (orally) on normal sleep, we used analysis of sleep architecture based on EEG and EMG recordings (Fig. 3A). Representative waveforms of EEG and EMG upon sleep states were shown in Fig. 3B. EEG power of delta (0.5–4 Hz) and theta (6–10 Hz) waves were measured by fast Fourier transformation (Fig. 3C). Figure 4 presents the effects of (–)- $\alpha$ -pinene (25, 50, and 100 mg/kg) and zolpidem (10 mg/kg) on sleep latency, NREMS, and REMS in C57BL/6N mice. Examples of EEG and EMG signals and corresponding hypnograms from a single mouse during the

first 3 hours were shown in Fig. 4A. Concentration of (–)- $\alpha$ -pinene was chosen from the pentobarbital-induced sleep test.

The values of sleep latency for (–)- $\alpha$ -pinene (100 mg/kg) and the positive control zolpidem (10 mg/kg) were  $19.6 \pm 3.5$  and



**Fig. 4.** Sleep-wake profiles in C57BL/6N mice after oral administration of (–)- $\alpha$ -pinene or zolpidem. (A) Typical signals of EEG and EMG recordings and the corresponding hypnograms in a mouse treated with vehicle, (–)- $\alpha$ -pinene, or zolpidem. (B) Effects of (–)- $\alpha$ -pinene and zolpidem on sleep latency. (C) Amount of NREMS and REMS during the 3-hour period after administration of vehicle, (–)- $\alpha$ -pinene, and zolpidem. Open and filled bars indicate the baseline day (vehicle administration) and experimental day [(–)- $\alpha$ -pinene or zolpidem administration], respectively. Each value represents the mean  $\pm$  S.E.M. of 8 mice in each group. \* $P < 0.05$ , \*\* $P < 0.01$ , significantly different from their vehicle (unpaired Student's *t* test).

8.1 ± 2.0 minutes, respectively (Fig. 4B). Both (–)- $\alpha$ -pinene ( $P < 0.05$ ) and zolpidem ( $P < 0.01$ ) produced a significant decrease in sleep latency. The decrease in sleep latency in mice treated with (–)- $\alpha$ -pinene indicates that it accelerates the initiation of NREMS just as zolpidem does. We calculated the amounts of NREMS and REMS during the first 3 hours after the administrations of (–)- $\alpha$ -pinene and zolpidem (Fig. 4C). As expected, the positive control zolpidem (10 mg/kg) increased the amount of NREMS by 1.7-fold ( $P < 0.01$ ) compared with that of the vehicle. Administration of (–)- $\alpha$ -pinene (100 mg/kg) was found to significantly increase the NREMS by 1.3-fold ( $P < 0.05$ ). There was no significant difference in the amount of REMS between (–)- $\alpha$ -pinene and zolpidem. Notably, in the pentobarbital-induced sleep test, (–)- $\alpha$ -pinene at 50 mg/kg significantly ( $P < 0.01$ ) decreased sleep latency and increased sleep duration (Fig. 2, A and B), whereas in the EEG and EMG experiment, (–)- $\alpha$ -pinene at the same dose did not show any significant difference (Fig. 4, B and C).


### Time Course Change of NREMS, REMS, and Wake

Figure 5 shows the time course of NREMS, REMS, and Wake for 24 hours after the administration of (–)- $\alpha$ -pinene (100 mg/kg) and zolpidem (10 mg/kg). (–)- $\alpha$ -Pinene (100 mg/kg) significantly increased the amount of NREMS during the second and third hours after administration by 1.37- and 1.58-fold relative to vehicle, respectively (Fig. 5A). This enhancement of NREMS was accompanied by a significant decrease in Wake during the same hours (Fig. 5A). The significant increase in NREMS by zolpidem (10 mg/kg) lasted for 5 hours after administration (Fig. 5B). Unlike zolpidem,


(–)- $\alpha$ -pinene did not produce a significant increase in NREMS for the first hour. Remarkably, both (–)- $\alpha$ -pinene and zolpidem did not affect REMS for 24 hours. After an initial increase in NREMS for the first 3 hours, there was no further significant change in sleep architecture during the subsequent periods (Fig. 5). This result indicates that (–)- $\alpha$ -pinene induced NREMS without causing adverse effects after sleep induction, which is consistent with previous findings (Masaki et al., 2012).

### Characteristics of Sleep-wake Episodes and Power Density

To better understand the nature of the sleep-enhancing effects of (–)- $\alpha$ -pinene, we additionally analyzed the total number and mean duration of NREMS, REMS, and Wake



**Fig. 5.** Time course changes in NREMS, REMS, and Wake in C57BL/6N mice after oral administration of (–)- $\alpha$ -pinene (A) and zolpidem (B) during the 24-hour period. Open and filled circles indicate the baseline day (vehicle administration) and experimental day [(–)- $\alpha$ -pinene or zolpidem administration], respectively. Each circle represents the hourly mean ± S.E.M. amount of each stage ( $n = 8$ ). \* $P < 0.05$ , \*\* $P < 0.01$ , significantly different from their vehicle (unpaired Student's  $t$  test). The horizontal filled and open bars on the x-axis (clock time) indicate the 12-hour dark and 12-hour light periods, respectively. Vehicle, (–)- $\alpha$ -pinene, and zolpidem were administered at 17:00.



**Fig. 6.** Characteristics of sleep-wake episodes in C57BL/6N mice after oral administration of (–)- $\alpha$ -pinene (A) and zolpidem (B). (A) Total number and mean duration of NREMS, REMS, and Wake bouts for 3 hours after the administration of (–)- $\alpha$ -pinene and zolpidem. (B) Sleep-wake stage transitions during the 3-hour period after the administration of (–)- $\alpha$ -pinene and zolpidem. Open and filled bars indicate the baseline day (vehicle administration) and experimental day [(–)- $\alpha$ -pinene or zolpidem administration], respectively. Each value represents the mean ± S.E.M. of 8 mice in each group. (C) EEG power density curves of NREMS caused by (–)- $\alpha$ -pinene and zolpidem. Delta activity, an index of sleep intensity, is shown in the inset histogram. The bar (—) represents the range of the delta wave (0.5–4 Hz). \* $P < 0.05$ , \*\* $P < 0.01$ , significantly different from their vehicle (unpaired Student's  $t$  test). N, non-rapid eye movement sleep; R, rapid eye movement sleep; W, wakefulness.

episodes, as well as sleep stage transition and EEG power density (Fig. 6). Both  $(-)\alpha$ -pinene and zolpidem significantly increased the number of bouts of Wake [ $(-)\alpha$ -pinene: 1.9-fold,  $P < 0.05$ ; zolpidem: 2.1-fold,  $P < 0.01$ ] and NREMS [ $(-)\alpha$ -pinene: 40%,  $P < 0.05$ ; zolpidem: 70%,  $P < 0.01$ ] but not REMS (Fig. 6A). Moreover, both  $(-)\alpha$ -pinene and zolpidem significantly decreased the duration of Wake [ $(-)\alpha$ -pinene: 52%,  $P < 0.05$ ; zolpidem: 56.5%,  $P < 0.01$ ] without affecting that of NREMS or REMS (Fig. 6A). Finally, both  $(-)\alpha$ -pinene and zolpidem significantly increased the number of state transitions from Wake to NREMS and from NREMS to Wake, whereas the number of stage transitions from NREMS to REMS and from REMS to NREMS were not affected. These results indicate that  $(-)\alpha$ -pinene inhibited the maintenance of Wake, consistent with the previous report (Omori et al., 2012).


So far, we observed an enhancement of quantity of sleep by  $(-)\alpha$ -pinene as evidenced by decrease of sleep latency and increase of NREMS time. Next, to evaluate the sleep intensity, we analyzed the EEG power density in mice during NREMS and measured the delta activity. We found that  $(-)\alpha$ -pinene did not affect the EEG power density (0–20 Hz) including delta activity (frequency range of 0.5–4 Hz) in NREMS compared with vehicle (Fig. 6C), whereas zolpidem significantly decreased the delta activity, indicating a slight loss of sleep intensity by zolpidem (Fig. 6C). These results suggest that, unlike zolpidem,  $(-)\alpha$ -pinene increased the quantity of sleep without compromising the sleep intensity.


### Molecular Mechanism of $(-)\alpha$ -Pinene

It has been reported that some monoterpenes such as borneol, verbenol, and pinocarveol act as a positive modulator of the GABA<sub>A</sub> receptors (Granger et al., 2005; Kessler et al., 2014). For this reason, we first investigated whether the GABAergic system

is involved in the hypnotic effects of  $(-)\alpha$ -pinene. To confirm the molecular mechanism of  $(-)\alpha$ -pinene on GABA<sub>A</sub>-BZD receptors, flumazenil (the antagonist at the GABA<sub>A</sub>-BZD receptor) was pretreated 15 minutes before administration of  $(-)\alpha$ -pinene and then analyzed the properties of sleep. First, we found that injection of flumazenil (1 mg/kg) alone did not produce significant changes in the sleep architecture (Fig. 7A) and concluded that flumazenil is not an inhibitor of sleep. Second, the hypnotic effect of zolpidem, the well-known GABA<sub>A</sub>-BZD receptor agonist, was fully antagonized by flumazenil (Fig. 7A). Finally, flumazenil also completely inhibited the hypnotic effect of  $(-)\alpha$ -pinene (Fig. 7A). The time course plot of duration of each sleep stage showed that in the presence of flumazenil  $(-)\alpha$ -pinene failed to alter the sleep architecture during 24 hours (Fig. 7B). These results suggest that the sleep-enhancing effects of  $(-)\alpha$ -pinene could be due to its modulation of GABA<sub>A</sub> receptor by acting at the BZD site.

To test the effect of  $(-)\alpha$ -pinene on GABA<sub>A</sub>-BZD receptors at GABAergic synapses, we performed whole cell patch-clamp recordings of IPSCs from hippocampal CA1 pyramidal neurons. These neurons are known to contain  $\alpha 1$ - and  $\alpha 2$ -subunits of GABA<sub>A</sub> receptors, which are the known targets of BZD drugs (Wisden et al., 1992; Somogyi et al., 1996). We found that zolpidem (10  $\mu$ M) significantly enhanced the decay time constant of sIPSCs in dose-dependent manner (Fig. 8, B and C) without affecting the amplitude and frequency of sIPSCs (Fig. 8, D and E), consistent with previous reports (Perrais and Ropert, 1999; Woo et al., 2014). We found that  $(-)\alpha$ -pinene (10  $\mu$ M) also enhanced the decay time constant of sIPSCs in dose-dependent manner ( $EC_{50}$ : 4.9 nM, Fig. 8, G and H), without affecting the amplitude and frequency of sIPSCs (Fig. 8, I and J). This enhancement of decay time constant by  $(-)\alpha$ -pinene was fully inhibited by flumazenil (1  $\mu$ M) (Fig. 8G). These results suggest that  $(-)\alpha$ -pinene prolongs the





**Fig. 8.** (A) Representative traces of sIPSC before (CON) and after treatment of zolpidem (10  $\mu$ M). (B) Averaged sIPSCs after normalization by peak (left). Decay was fitted to one-exponential functions. Summary bar graph of sIPSC decay value before and after treatment of zolpidem (right). Data are represented as mean  $\pm$  S.E.M. \*\* $P < 0.01$ , Students' tailed  $t$  test. (C–E) Summary graphs of sIPSC decay tau value (C), amplitude (D), and frequency (E) after normalization by control response. (F) Representative traces of sIPSC before (CON) and after treatment of (-)- $\alpha$ -pinene (10  $\mu$ M). (G) Averaged sIPSCs after normalization by peak (left). Summary bar graph of sIPSC decay value before and after treatment of (-)- $\alpha$ -pinene and flumazenil (1  $\mu$ M, right). \* $P < 0.05$ , one-way ANOVA test. (H–J). Summary graphs of sIPSC decay tau value (H), amplitude (I), and frequency (J) after normalization by control response. Decay response was fitted using sigmoidal logistic 4 parameters.

GABAergic synaptic transmission effectively by modulating the GABA<sub>A</sub> BZD receptor.


### Binding Mode of (-)- $\alpha$ -Pinene in BZD Binding Site of GABA<sub>A</sub> Receptor

To predict the binding modes of zolpidem, flumazenil and (-)- $\alpha$ -pinene at the BZD binding site, the human  $\alpha 1\beta 2\gamma 2$  subtype of GABA<sub>A</sub> receptor was generated based on the X-ray crystal structure of  $\beta 3$  homopentamer GABA<sub>A</sub> receptor (PDB: 4COF; Fig. 9A, Supplemental Fig. 1) (Miller and Aricescu, 2014). Each compound was docked into the BZD binding site between  $\alpha 1$ - and  $\gamma 2$ -subunits in extracellular domain of GABA<sub>A</sub> receptor using InduceFit docking method of Maestro Program. As shown in Fig. 9, B and C, and Supplemental Figs. 2 and 3, the binding mode of zolpidem and flumazenil commonly showed that: 1)  $\alpha 1\text{Tyr}209$  is engaged in  $\pi$ - $\pi$  interaction with phenyl rings and 2) amine groups of loop C  $\alpha 1\text{Ser}205$  are forming a hydrogen bond of carbonyl group of compounds. This is consistent with the fact that  $\alpha 1\text{Tyr}209$  is crucial for high binding affinity of diazepam, zolpidem, and flumazenil as shown by mutagenesis studies (Amin et al., 1997; Buhr et al., 1997). In addition, zolpidem participates three  $\pi$ - $\pi$  interactions with loop D  $\gamma 2\text{Phe}77$ , loop A  $\alpha 1\text{His}101$ , and loop D  $\gamma 2\text{Ala}79$ , whereas flumazenil has hydrophobic contacts with  $\gamma 2\text{Tyr}58$ ,  $\gamma 2\text{Phe}77$ , and  $\gamma 2\text{Met}130$ . This is consistent with experimental data that mutations of  $\gamma 2\text{Phe}77$  and  $\alpha 1\text{His}101$  lead to loss of the affinity for zolpidem, whereas the same mutations lead to

only slight change in the affinity for flumazenil (Wieland et al., 1992; Buhr et al., 1997). The binding energy of (-)- $\alpha$ -pinene with Gscore of -6.57 kcal/mol was lower than zolpidem and flumazenil with Gscore of -9.22 and -8.78 kcal/mol, respectively. These results predict that (-)- $\alpha$ -pinene probably show lower binding activities than zolpidem and flumazenil. The  $\alpha$ -pinene makes strong hydrophobic interactions with aromatic residues of  $\alpha 1\text{Phe}99$ ,  $\alpha 1\text{Tyr}159$ ,  $\alpha 1\text{Tyr}209$ , and  $\gamma 2\text{Phe}77$  (Fig. 9D and Supplemental Fig. 4). These results indicate that (-)- $\alpha$ -pinene may positively modulate the biologic function of GABA<sub>A</sub> receptor by directly binding at the BZD binding site

## Discussion

We demonstrated that (-)- $\alpha$ -pinene enhances NREMS by prolonging the decay time constant of GABAergic synaptic transmission by directly acting at BZD binding site of GABA<sub>A</sub> receptor. Until now, it has been reported that inhalation of (-)- $\alpha$ -pinene produces sedative and anxiolytic effects. To our best knowledge, this is the first study to investigate the effects of oral administration of (-)- $\alpha$ -pinene on sleep and its hypnotic mechanism. We report that the effect of (-)- $\alpha$ -pinene in sleep was specific in NREMS rather than REMS by oral administration in mice (Fig. 4). Contrary to our findings, it has been reported that there was an increase of REMS by inhaled (+)- $\alpha$ -pinene in rat (Satou et al., 2014). This discrepancy might be due to different administration method, concentration, enantiomer type, or species.



**Fig. 9.** Putative binding modes in BZD binding site of GABA<sub>A</sub> receptor. Top and side views of GABA<sub>A</sub> receptor homology model (A), the binding pose of zolpidem (B), flumazenil (C), and (–)- $\alpha$ -pinene (D) obtained by InduceFit docking method implemented in Schrödinger program. The PDB files of putative binding modes are in Supplemental Figs. 1–4. Representation shows the  $\alpha$ 1-subunit (blue),  $\beta$ 2-subunit (Green) and  $\gamma$ 2-subunit (red). Ligand and key residues are shown with stick. Hydrogen bond interactions are depicted by green dotted line and hydrophobic interactions are indicated by purple dotted line. Square denotes the BZD binding site between  $\alpha$ 1-subunit and  $\gamma$ 2-subunit in the human GABA<sub>A</sub> receptor. ECD, extracellular domain; TMD, transmembrane domain.

It has been reported that sleep quantity, as indicated by increased duration of NREMS, is enhanced by various hypnotics, including natural compounds as well as well-known drugs such as diazepam and zolpidem (Cho et al., 2011; Chen et al., 2012; Masaki et al., 2012). In addition to an increase in duration of NREMS, zolpidem can increase sleep quantity by changing the sleep architecture such as changes in bout number and sleep state transition (Chen et al., 2012; Qu et al., 2012). We found that, just like zolpidem, (–)- $\alpha$ -pinene not only enhanced the duration of NREMS (Fig. 4) but also decreased the Wake bout (Fig. 6) and increased the number of transition in Wake to NREMS and NREMS to Wake (Fig. 6). Based on these results, we conclude that (–)- $\alpha$ -pinene acts as a hypnotic by enhancing sleep quantity.

When it comes to “good sleep,” it is important to consider sleep quality in addition to sleep quantity. Although it is difficult to define sleep quality in rodents, it has been widely accepted that delta (0.5–4 Hz) activity could be a good indicator of the quality or intensity of NREMS (Tobler et al., 2001; Chen et al., 2012). In human and rodent, it has been reported that diazepam and zolpidem increase the sleep quantity in NREMS but paradoxically reduce the delta activity (Feinberg et al., 2000; Tobler et al., 2001). In our study, zolpidem produced a typical decrease in delta activity as expected. Interestingly, (–)- $\alpha$ -pinene did not alter delta activity (Fig. 6C), consistent with other natural compounds that show hypnotic effects (Cho et al., 2014; Yoon et al., 2014). Therefore, (–)- $\alpha$ -pinene shows an advantage over zolpidem or diazepam in that it preserves intensity of NREMS.

A bicyclic monoterpene, borneol, which is found in essential oils for analgesia and anesthesia in traditional Chinese medicine, acts as a positive modulator at GABA<sub>A</sub> receptors (Granger et al., 2005). It has been reported that the other monoterpenes, verbenol and pinocarveol, potentiated the GABA action of the GABA<sub>A</sub> receptors (Kessler et al., 2014). These reports suggest that hypnotic effect of (–)- $\alpha$ -pinene may involve GABAergic mechanism. Therefore, to demonstrate the

action mechanism of (–)- $\alpha$ -pinene, the effect of flumazenil on the hypnotic effect of (–)- $\alpha$ -pinene was tested. Flumazenil is a well-known antagonist of the GABA<sub>A</sub>-BZD receptors and inhibits the effects of BZD agonists such as zolpidem and diazepam by competitively blocking their binding (Johnston, 2005; Chen et al., 2012). Hypnotic effects of zolpidem and (–)- $\alpha$ -pinene were completely blocked by flumazenil (Fig. 7). These findings support that the hypnotic effects of (–)- $\alpha$ -pinene should be attributed to the positive allosteric modulation of GABA<sub>A</sub> receptors at the BZD-binding site via a mechanism similar to that of zolpidem.

BZD agonists such as diazepam and zolpidem are known to enhance GABAergic inhibitory signaling by prolonging the decay time constant of GABA<sub>A</sub> receptors in various brain regions including thalamus, hippocampus, and neocortex (Perrais and Ropert, 1999; Bacci et al., 2003; Woo et al., 2014). In our recent study, we reported that the single compound, isoliquiritigenin, derived from flavonoids shows the same effects of prolonging the decay time constant of GABA<sub>A</sub> receptor-mediated IPSCs, acting as a modulator of GABA<sub>A</sub>-BZD receptors and thus enhancing the quantity of NREMS (Cho et al., 2011; Woo et al., 2014). In the current study, we found the same effects of (–)- $\alpha$ -pinene. On the basis

**TABLE 1**  
Comparison of effectiveness between zolpidem and (–)- $\alpha$ -pinene in electrophysiology, modeling, and sleep behavior

	Zolpidem	(–)- $\alpha$ -Pinene
Electrophysiology		
Affinity	Low	High
Efficacy	High	Low
Potency (at 1 $\mu$ M)	High	Low
Molecular modeling		
Binding energy	High	Low
Sleep effect		
Quantity	Increase	Increase
Intensity	Decrease	No effect



of these findings, we can predict based on the ability to prolong the decay time constant of IPSCs whether a certain drug or chemical can act as a potential modulator acting at the BZD site of GABA<sub>A</sub>-BZD receptors and possibly enhance the quantity of NREMS. This assay can serve as a simple screening method for finding potential hypnotics mimicking benzodiazepines.

The molecular modeling results demonstrate that (–)- $\alpha$ -pinene directly binds to the BZD binding site of GABA<sub>A</sub>. To investigate the binding mode of (–)- $\alpha$ -pinene at the BZD binding site of GABA<sub>A</sub>, we generated a homology model of  $\alpha$ 1 $\beta$ 2 $\gamma$ 2 GABA<sub>A</sub> receptor, which is most abundant in the physiologic system, based on the X-ray crystal structure of  $\beta$ 3 homopentamer GABA<sub>A</sub> receptor (Miller and Aricescu, 2014), because the  $\beta$ 3 homopentamer lacks the BZD binding site. We conducted molecular docking study of (–)- $\alpha$ -pinene including zolpidem and flumazenil at the BZD binding site of putative model of  $\alpha$ 1 $\beta$ 2 $\gamma$ 2 GABA<sub>A</sub> receptor. The binding modes of zolpidem and flumazenil at the BZD binding site between  $\alpha$ 1- and  $\gamma$ 2-subunits were in good agreement with experimental study of mutagenesis. It has been reported that the residue of  $\alpha$ 1Tyr209 is essential for binding of ligands at BZD binding site, whereas the residues of  $\alpha$ 1His101 and  $\gamma$ 2Phe77 are needed for binding of zolpidem but not for flumazenil (Wieland et al., 1992; Buhr et al., 1997). The (–)- $\alpha$ -pinene was well accommodated into the binding pocket and shared the similar hydrophobic interactions as flumazenil and zolpidem, with key residues of  $\alpha$ 1Phe99,  $\alpha$ 1Tyr159,  $\alpha$ 1Tyr209, and  $\gamma$ 2Phe77 at the BZD binding site.

From the molecular modeling, we obtained binding energy for each ligand and its rank order: (–)- $\alpha$ -pinene (–6.57 kcal/mol) < flumazenil (–8.78 kcal/mol) < zolpidem (–9.22 kcal/mol). Based on this rank order, we can make various predictions on physiologic effects of each ligand. First of all, it is possible that the deficiency of both  $\pi$ - $\pi$  interaction of (–)- $\alpha$ -pinene with  $\alpha$ 1Tyr209 and hydrogen bond of (–)- $\alpha$ -pinene with  $\alpha$ 1Ser205 might have resulted in lower binding energy, lower potency, and lower efficacy than those of zolpidem and flumazenil. For example, zolpidem at 1  $\mu$ M enhanced the sIPSC decay by about 35%, whereas (–)- $\alpha$ -pinene enhanced by about 20% at the same concentration, indicating a lower potency of (–)- $\alpha$ -pinene than zolpidem. The efficacy of (–)- $\alpha$ -pinene, which at 1  $\mu$ M already showed saturation at 30% of decay, appears to be lower than zolpidem, which has not been saturated even at 10  $\mu$ M (about 60%), suggesting that (–)- $\alpha$ -pinene might act as a partial modulator at the BZD binding site of GABA<sub>A</sub> receptor. These rank orders of potency and efficacy are in line with the rank order of binding energy for (–)- $\alpha$ -pinene and zolpidem. In contrast, (–)- $\alpha$ -pinene showed higher affinity (EC<sub>50</sub>: 4.9 nM) than zolpidem in sIPSC decay, which did not correlate well with the rank order of binding energy.

In summary, when we compare the effectiveness between zolpidem and (–)- $\alpha$ -pinene based on electrophysiology, molecular modeling, and sleep behavior, the sleep quantity was positively correlated with efficacy, potency, and binding energy, whereas the sleep intensity was not (Table 1). Furthermore, the sleep intensity was preserved when (–)- $\alpha$ -pinene acted as a partial modulator. Although (–)- $\alpha$ -pinene showed lower binding energy and efficacy than zolpidem, the effectiveness of (–)- $\alpha$ -pinene for sleep seems to be better than zolpidem when we consider both sleep quantity and sleep intensity. Based on these results, we propose that various parameters obtained from electrophysiology and molecular modeling (e.g., affinity, efficacy, potency, and binding energy) could be a good marker for screening of novel drugs for sleep.

In conclusion, by combining sleep behavior analysis, electrophysiology, and molecular modeling, we demonstrate the feasibility of finding a sleep modulator. We report that (–)- $\alpha$ -pinene can be a useful hypnotic by its potent action at the BZD site of GABA<sub>A</sub> receptors as well as by its easy accessibility. In addition to hypnotic role of (–)- $\alpha$ -pinene, it has been shown to display a variety of beneficial properties such as anxiolytic, anti-inflammatory, antioxidant. Most importantly, we can obtain these beneficial effects of (–)- $\alpha$ -pinene from our daily life (e.g., strolling in woods or inhaling essential oil). Through our demonstration on the hypnotic effect of (–)- $\alpha$ -pinene by oral administration, we propose that (–)- $\alpha$ -pinene could be a good therapeutic agent for treating sleeping disorder or anxiety.

#### Authorship Contributions

Participated in research design: Yang, Woo, Lee, and Cho.

Conducted experiments: Yang, Woo, Um, and Yoon.

Contributed new reagents of analytic tools: Pae, Park, and Cho.

Wrote or contributed to the writing of the manuscript: Yang, Woo, Pae, Um, N.-C. Cho, Lee, and S. Cho.

#### References

- Amin J, Brooks-Kayal A, and Weiss DS (1997) Two tyrosine residues on the alpha subunit are crucial for benzodiazepine binding and allosteric modulation of gamma-aminobutyric acid A receptors. *Mol Pharmacol* **51**:833–841.
- Bacci A, Rudolph U, Huguenard JR, and Prince DA (2003) Major differences in inhibitory synaptic transmission onto two neocortical interneuron subclasses. *J Neurosci* **23**:9664–9674.
- Bakkali F, Averbeck S, Averbeck D, and Idaomar M (2008) Biological effects of essential oils—a review. *Food Chem Toxicol* **46**:446–475.
- Buhr A, Baur R, and Sigel E (1997) Subtle changes in residue 77 of the gamma subunit of alpha1beta2gamma2 GABA<sub>A</sub> receptors drastically alter the affinity for ligands of the benzodiazepine binding site. *J Biol Chem* **272**:11799–11804.
- Chen CR, Zhou XZ, Luo YJ, Huang ZL, Urade Y, and Qu WM (2012) Magnolol, a major bioactive constituent of the bark of *Magnolia officinalis*, induces sleep via the benzodiazepine site of GABA(A) receptor in mice. *Neuropharmacology* **63**:1191–1199.
- Cho S, Kim S, Jin Z, Yang H, Han D, Baek NI, Jo J, Cho CW, Park JH, Shimizu M, et al. (2011) Isoliquiritigenin, a chalcone compound, is a positive allosteric modulator of GABA<sub>A</sub> receptors and shows hypnotic effects. *Biochem Biophys Res Commun* **413**:637–642.
- Cho S, Yoon M, Pae AN, Jin YH, Cho NC, Takata Y, Urade Y, Kim S, Kim JS, Yang H, et al. (2014) Marine polyphenol phlorotannins promote non-rapid eye movement sleep in mice via the benzodiazepine site of the GABA<sub>A</sub> receptor. *Psychopharmacology (Berl)* **231**:2825–2837.
- FDA (2015) Code of Federal Regulations Title 21. Food and Drug Administration, Washington, DC.
- Feinberg I, Maloney T, and Campbell IG (2000) Effects of hypnotics on the sleep EEG of healthy young adults: new data and psychopharmacological implications. *J Psychiatr Res* **34**:423–438.
- Gomes-Carneiro MR, Viana ME, Felzenszwalb I, and Paumgarten FJ (2005) Evaluation of beta-myrcene, alpha-terpinene and (+)- and (-)-alpha-pinene in the Salmonella/microsome assay. *Food Chem Toxicol* **43**:247–252.
- Granger RE, Campbell EL, and Johnston GAR (2005) (+)- And (-)-borneol: efficacious positive modulators of GABA action at human recombinant alpha1beta2gamma2L GABA(A) receptors. *Biochem Pharmacol* **69**:1101–1111.
- Groot PDG and MacDonald L (2002) Influence of enantiomers of  $\alpha$ -pinene on the response of the red pine cone beetle, *Conophthorus resinosae* to its pheromone pinyol. *Entomol Exp Appl* **105**:169–174.
- Him A, Ozbek H, Turel I, and Oner AC (2008) Antinociceptive activity of alphapinene and fenchone. *Pharmacology Online* **3**:363–369.
- Hmamouchi M, Hamamouchi J, Zouhdi M, and Bessiere JM (2001) Chemical and Antimicrobial Properties of Essential Oils of Five Moroccan Pinaceae. *J Essent Oil Res* **13**:298–302.
- Inoannou E, Koutsaviti A, Tzakou O, and Roussis V (2014) The genus *Pinus*: a comparative study on the needle essential oil composition of 46 pine species. *Phytochem Res* **13**:741–768.
- Johnston GA (2005) GABA(A) receptor channel pharmacology. *Curr Pharm Des* **11**:1867–1885.
- Judzientiene A and Kupcinskiene E (2008) Chemical composition on essential oils from needles of *Pinus sylvestris* L. grown in northern Lithuania. *J Essent Oil Res* **20**:26–29.
- Kamal EH, Al-Ajmi MF, and Abdullah Ma-B (2003) Some cardiovascular effects of the dethymoquinonated *Nigella sativa* volatile oil and its major components  $\alpha$ -pinene and p-cymene in rats. *Saudi Pharm J* **11**:104–110.
- Kessler A, Sahin-Nadeem H, Lummis SCR, Weigel I, Pischetsrieder M, Buettner A, and Villmann C (2014) GABA(A) receptor modulation by terpenoids from *Sideritis* extracts. *Mol Nutr Food Res* **58**:851–862.
- Lee JG, Lee CG, Kwag JJ, Buglass AJ, and Lee GH (2005) Determination of optimum conditions for the analysis of volatile components in pine needles by double-shot pyrolysis-gas chromatography-mass spectrometry. *J Chromatogr A* **1089**:227–234.

- Limberger RP, Aleixo AM, Fett-Neto AG, and Henriques AT (2007) Bioconversion of (+)- and (-)-alpha pinene to (+)- and (-)-verbenone by plant cell cultures of *Psychotria brachyceras* and *Rauvolfia sellowii*. *Electron J Biotechnol* **10**:500–507.
- Masaki M, Aritake K, Tanaka H, Shoyama Y, Huang ZL, and Urade Y (2012) Crocin promotes non-rapid eye movement sleep in mice. *Mol Nutr Food Res* **56**:304–308.
- Miller PS and Aricescu AR (2014) Crystal structure of a human GABAA receptor. *Nature* **512**:270–275.
- Omori K, Kagami Y, Yokoyama C, Moriyama T, Matsumoto N, Masaki M, Nakamura H, Kamasaka H, Shiraishi K, Kometani T, et al. (2012) Promotion of non-rapid eye movement sleep in mice after oral administration of ornithine. *Sleep Biol Rhythms* **10**:38–45.
- Orhan I, K peli E, Aslan M, Kartal M, and Yesilada E (2006) Bioassay-guided evaluation of anti-inflammatory and antinociceptive activities of pistachio, *Pistacia vera* L. *J Ethnopharmacol* **105**:235–240.
- Perrais D and Ropert N (1999) Effect of zolpidem on miniature IPSCs and occupancy of postsynaptic GABAA receptors in central synapses. *J Neurosci* **19**:578–588.
- Qu WM, Yue XF, Sun Y, Fan K, Chen CR, Hou YP, Urade Y, and Huang ZL (2012) Honokiol promotes non-rapid eye movement sleep via the benzodiazepine site of the GABA(A) receptor in mice. *Br J Pharmacol* **167**:587–598.
- Rivas dSA, Lopes P, Barros dAM, Costa D, Alviano C, and Alviano D (2012) Biological activities of  $\alpha$ -pinene and  $\beta$ -pinene enantiomers. *Molecules* **17**: 6305–6316.
- Sali A and Blundell TL (1993) Comparative protein modelling by satisfaction of spatial restraints. *J Mol Biol* **234**:779–815.
- Satou T, Kasuya H, Maeda K, and Koike K (2014) Daily inhalation of  $\alpha$ -pinene in mice: effects on behavior and organ accumulation. *Phytother Res* **28**:1284–1287.
- Somogyi P, Fritschy JM, Benke D, Roberts JDB, and Sieghart W (1996) The gamma 2 subunit of the GABAA receptor is concentrated in synaptic junctions containing the alpha 1 and beta 2/3 subunits in hippocampus, cerebellum and globus pallidus. *Neuropharmacology* **35**:1425–1444.
- S ntar I, Tumen I, Ust n O, Keleş H, and Akkol EK (2012) Appraisal on the wound healing and anti-inflammatory activities of the essential oils obtained from the cones and needles of *Pinus* species by in vivo and in vitro experimental models. *J Ethnopharmacol* **139**:533–540.
- Tobler I, Kopp C, Deboer T, and Rudolph U (2001) Diazepam-induced changes in sleep: role of the alpha 1 GABA(A) receptor subtype. *Proc Natl Acad Sci USA* **98**: 6464–6469.
- Wieland HA, L ddens H, and Seeburg PH (1992) A single histidine in GABAA receptors is essential for benzodiazepine agonist binding. *J Biol Chem* **267**: 1426–1429.
- Wisden W, Laurie DJ, Monyer H, and Seeburg PH (1992) The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci* **12**:1040–1062.
- Woo J, Cho S, and Lee CJ (2014) Isoliquiritigenin, a chalcone compound, enhances spontaneous inhibitory postsynaptic response. *Exp Neurobiol* **23**:163–168.
- Xie Q, Liu Z, and Li Z (2015) Chemical composition and antioxidant activity of essential oil of six *Pinus* taxa native to China. *Molecules* **20**:9380–9392.
- Yamaoka S, Tomita T, Imaizumi Y, Watanabe K, and Hatanaka A (2005) Effects of plant-derived odors on sleep-wakefulness and circadian rhythmicity in rats. *Chem Senses* **30** (Suppl 1):i264–i265.
- Yang X, Zhao HT, Wang J, Meng Q, Zhang H, Yao L, Zhang YC, Dong AJ, Ma Y, Wang ZY, et al. (2010) Chemical composition and antioxidant activity of essential oil of pine cones of *Pinus armandii* from the Southwest region of China. *J Med Plants Res* **4**:1668–1672.
- Yoon M, Kim JS, Jo J, Han D, and Cho S (2014) Sleep-promoting effect of *Ecklonia* cava: Ethanol extract promotes non-rapid eye movement sleep in C57BL/6N Mice. *Fish Aqua Sci* **17**: 19–25.

---

**Address correspondence to:** Dr. C. Justin Lee, Director of Center for Neural Science, Korean Institute of Science and Technology, 5, Hwarang-ro 14-gil, Seoungbuk-gu, Seoul 02792, Republic of Korea. E-mail: cjl@kist.re.kr or Dr. Suengmok Cho, Principal Research Scientist, Professional Engineer for Food, Functional Food Laboratory for Sleep Regulation, Korea Food Research Institute, 62, Anyangpangyo-ro 1201, Bundang-gu, Seongnam-si 13539, Republic of Korea. E-mail: smcho@kfri.re.kr

---