

# Elongation Factor 2 and Fragile X Mental Retardation Protein Control the Dynamic Translation of Arc/Arg3.1 Essential for mGluR-LTD

Sungjin Park,<sup>1,7</sup> Joo Min Park,<sup>1,7</sup> Sangmok Kim,<sup>1</sup> Jin-Ah Kim,<sup>1</sup> Jason D. Shepherd,<sup>4</sup> Constance L. Smith-Hicks,<sup>2</sup> Shoaib Chowdhury,<sup>1</sup> Walter Kaufmann,<sup>2</sup> Dietmar Kuhl,<sup>5</sup> Alexey G. Ryazanov,<sup>6</sup> Richard L. Huganir,<sup>1,3</sup> David J. Linden,<sup>1</sup> and Paul F. Worley<sup>1,2,\*</sup>

<sup>1</sup>Department of Neuroscience

<sup>2</sup>Department of Neurology

<sup>3</sup>Howard Hughes Medical Institute

Johns Hopkins University School of Medicine, Baltimore, MD 20205, USA

<sup>4</sup>The Picower Institute for Learning and Memory, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

<sup>5</sup>Molecular Neurobiology, Department of Biology-Chemistry-Pharmacy, Freie Universität Berlin, 14195 Berlin, Germany

<sup>6</sup>Department of Pharmacology, University of Medicine and Dentistry of New Jersey-Robert Wood Johnson Medical School, Piscataway, NY 08854, USA

<sup>7</sup>These authors contributed equally to this work

\*Correspondence: [pworley@jhmi.edu](mailto:pworley@jhmi.edu)

DOI 10.1016/j.neuron.2008.05.023

## SUMMARY

Group I metabotropic glutamate receptors (mGluR) induce long-term depression (LTD) that requires protein synthesis. Here, we demonstrate that Arc/Arg3.1 is translationally induced within 5 min of mGluR activation, and this response is essential for mGluR-dependent LTD. The increase in Arc/Arg3.1 translation requires eEF2K, a Ca<sup>2+</sup>/calmodulin-dependent kinase that binds mGluR and dissociates upon mGluR activation, whereupon it phosphorylates eEF2. Phospho-eEF2 acts to slow the elongation step of translation and inhibits general protein synthesis but simultaneously increases Arc/Arg3.1 translation. Genetic deletion of *eEF2K* results in a selective deficit of rapid mGluR-dependent Arc/Arg3.1 translation and mGluR-LTD. This rapid translational mechanism is disrupted in the fragile X disease mouse (*Fmr1* KO) in which mGluR-LTD does not require de novo protein synthesis but does require Arc/Arg3.1. We propose a model in which eEF2K-eEF2 and FMRP coordinately control the dynamic translation of Arc/Arg3.1 mRNA in dendrites that is critical for synapse-specific LTD.

## INTRODUCTION

Long-lasting forms of synaptic plasticity require de novo protein synthesis (Kelleher et al., 2004; Wang and Tiedge, 2004). N-methyl-D-aspartate (NMDA) receptor-dependent long-term potentiation (LTP) of the Schaffer collateral-CA1 synapse lasting longer than ~60 min is blocked by agents that halt translation. Late-phase NMDA-dependent LTD in hippocampal slice cultures is also protein synthesis dependent (Kauderer and Kandel,

2000). De novo protein synthesis is also required for forms of long-term depression (LTD) that are induced by group I metabotropic glutamate receptor (mGluR) activation or by paired-pulse low-frequency stimulation (PP-LFS) (Huber et al., 2000). In contrast to NMDA receptor-dependent LTP and LTD, where the requirement for protein synthesis is delayed, mGluR-LTD requires de novo protein synthesis within 5–10 min. (Huber et al., 2000).

The products of de novo protein synthesis are hypothesized to be “captured” at active synapses (Frey and Morris, 1997), but their identity remains unknown. Arc/Arg3.1 is an immediate-early gene (IEG) that is induced by NMDA receptor activation in vivo (Link et al., 1995; Lyford et al., 1995) and mediates a postsynaptic endocytic pathway by interacting with endophilin 2/3 and dynamin that selectively traffics AMPA receptors (AMPA) (Chowdhury et al., 2006). The activity-dependent expression of Arc/Arg3.1 mRNA and protein underlie a homeostatic mechanism that maintains a precise level of AMPAR-dependent excitability in conditions of persistently increased or decreased synaptic input (Shepherd et al., 2006). Studies presented here demonstrate that Arc/Arg3.1 is also required for mGluR and PP-LFS LTD. In contrast to changes in Arc/Arg3.1 expression that occur over hours to days in homeostatic plasticity, mGluR activation results in increases in Arc/Arg3.1 protein within 3–5 min. Using biochemical and genetic approaches, we demonstrate that mGluR evokes rapid Arc/Arg3.1 translation via a signaling pathway that involves eukaryotic elongation factor 2 kinase (eEF2K) and eukaryotic elongation factor 2 (eEF2). eEF2 is required for the elongation step of translation (Ryazanov et al., 1988), while phospho-eEF2 acts as a potent inhibitor of the elongation step in a manner that is similar to chemical protein synthesis inhibitors, such as cycloheximide (Begueret et al., 1977; Obrig et al., 1971). Our data support a model in which rapid translational upregulation of Arc/Arg3.1 is required for mGluR-dependent LTD, and this translational induction is generated as a consequence of local and transient inhibition of the translation of other mRNAs via phospho-eEF2.

Our studies also provide insights into the molecular basis of altered synaptic plasticity in fragile X mental retardation syndrome. Fragile X syndrome, which is the most common inherited cause of mental retardation and autism (O'Donnell and Warren, 2002), is caused by an expansion of CGG in the 5' untranslated region of the fragile X mental retardation protein (FMRP) gene (*Fmr1*) that reduces its expression. FMRP binds to G-quartet-containing RNAs through the RGG box (Darnell et al., 2001; Schaeffer et al., 2001). FMRP functions as a translational repressor of specific synaptic mRNAs by a proposed mechanism that may involve BC1 RNA, a nontranslatable message abundant in dendrites (Zalfa et al., 2003; but see also Iacoangeli et al., 2008). FMRP is also reported to function as part of a RISC nuclease complex that represses translation by RNA interference (Caudy et al., 2002; Ishizuka et al., 2002; Jin et al., 2004). In *Fmr1* knockout (KO) mice, Arc/Arg3.1,  $\alpha$ -CaMKII, and MAP1B proteins are reported to be elevated both in total brain and synaptosomal fractions (Zalfa et al., 2003). Notably, an increased association of Arc/Arg3.1 mRNA with polyribosomes suggests that a greater fraction of Arc/Arg3.1 mRNA is being actively translated in *Fmr1* KO. The notion that misregulated expression of these proteins might underlie cognitive deficits in fragile X syndrome is supported by physiological studies that implicate FMRP in altered synaptic plasticity. *Fmr1* KO mice show robust mGluR-LTD (Huber et al., 2002) that is insensitive to protein synthesis inhibitors (Hou et al., 2006; Nosyreva and Huber, 2006), suggesting that, when translation is derepressed due to the absence of FMRP, "LTD proteins" that are normally produced in a stimulus-dependent manner are now continuously synthesized and are sufficient to sustain mGluR-LTD without the requirement for de novo protein synthesis. Insight into the molecular basis of mGluR and FMRP-dependent translation is provided by the observations that the efficacy of FMRP to inhibit translation is regulated by phosphorylation (Ceman et al., 2003) and that mGluR activation results in dephosphorylation of FMRP by PP2A within 1 min that is linked to rapid translational upregulation of a target protein, SAPAP3 (Narayanan et al., 2007). A similar rapid mGluR- and FMRP-dependent increase in  $\alpha$ -CaMKII, PSD-95 (Muddashetty et al., 2007), and MAP1B (Davidkova and Carroll, 2007) has been reported, and MAP1B is suggested to play a role in glutamate receptor trafficking in cultured neurons. In the present study, we find that rapid, de novo Arc/Arg3.1 expression is disrupted in *Fmr1* KO neurons, and Arc/Arg3.1/*Fmr1* double KO mice show reduced mGluR-LTD. These observations suggest that eEF2K/eEF2 and FMRP-dependent translational pathways are cofunctional in controlling a rapid and transient switch of the translational machinery to mRNAs that are essential for mGluR-dependent synaptic depression.

## RESULTS

### mGluR-LTD and PP-LFS LTD Require Arc/Arg3.1

To examine the role of Arc/Arg3.1 in mGluR-LTD, we monitored Schaffer collateral-CA1 synapses in acute hippocampal slices prepared from wild-type (WT) and Arc/Arg3.1 knockout (KO) mice. Baseline synaptic properties, including the fiber volley-fEPSP relationship (an index of basal synaptic strength) and


paired-pulse facilitation ratio were normal in Arc/Arg3.1 KO mice (see Figures S1A and S1B available online), confirming a previous report (Plath et al., 2006). In WT slices, treatment with the group I mGluR agonist DHPG (50  $\mu$ M) for 5 min followed by washout produced a stable reduction of synaptic strength (72.8%  $\pm$  2.0% of baseline, mean  $\pm$  SEM) (Figure 1A). Synaptic stimulation using the paired-pulse low-frequency stimulation (PP-LFS) protocol in the presence of the NMDA receptor antagonist D-APV (50  $\mu$ M) resulted in a similar stable reduction of synaptic strength to 79.9%  $\pm$  2.1% of baseline (Figure 1B). In Arc/Arg3.1 KO slices, treatment with DHPG (92.1%  $\pm$  3.7% of baseline,  $p < 0.001$  compared to littermate WT controls by unpaired two-tailed Student's *t* test) or PP-LFS (94.3%  $\pm$  2.1% of baseline,  $p < 0.0001$ ) failed to evoke robust LTD, albeit there is a slight residual LTD in Arc/Arg3.1 KO slices ( $p = 0.09$  for DHPG-LTD and  $p = 0.03$  for PP-LFS LTD by paired *t* test). The residual LTD suggests that an Arc/Arg3.1-independent pathway also contributes to mGluR-LTD. The immediate short-term synaptic depression during the induction period with DHPG and immediately following the PP-LFS protocol was not significantly different between WT and KO mice (Figures 1A and 1B). Furthermore, mGluR-LTD induced by higher concentration of DHPG (100  $\mu$ M) was also impaired in Arc/Arg3.1 KO slices (88.5%  $\pm$  7.4% of baseline for Arc/Arg3.1 KO slices; 64.4%  $\pm$  2.6% of baseline for WT slices,  $p < 0.01$ ), indicating that the requirement for Arc/Arg3.1 does not depend on a specific range of mGluR activation (Figure S1C).

### mGluR-Dependent AMPA Receptor Endocytosis Requires Arc/Arg3.1

mGluR1/5 activation results in a rapid and sustained loss of surface AMPARs that underlies synaptic depression (Snyder et al., 2001). Since Arc/Arg3.1 KO mice have deficient mGluR-LTD, we directly tested whether Arc/Arg3.1 is required for mGluR-dependent AMPAR endocytosis. DHPG (50  $\mu$ M) was applied to DIV 14–21 primary hippocampal neurons for 5 min followed by washout, and surface and internalized AMPARs were measured 15 or 60 min after DHPG application. In WT cultures, DHPG resulted in a significant loss of surface GluR1 at 15 and 60 min as compared with untreated cultures (Figures 1C1–1C3 and 1C7). However, GluR1 surface levels were unchanged after DHPG application in Arc/Arg3.1 KO neurons (Figures 1C4–1C6 and 1C7). WT cultures exhibited a significant increase in internalized GluR1 at 15 min (Figures 1D1–1D3 and 1D7). Arc/Arg3.1 KO neurons did not exhibit an increase in internalized receptors after DHPG application (Figures 1D4–1D6 and 1D7). Thus, we conclude that Arc/Arg3.1 is required for rapid, mGluR-dependent AMPAR endocytosis.

### mGluR Induces Rapid Translation of Preexisting Arc/Arg3.1 mRNA

If Arc/Arg3.1 plays a direct role in mGluR-LTD, we anticipated that its protein level should be acutely regulated in dendrites. Therefore, Arc/Arg3.1 protein expression was examined by immunocytochemistry in DIV14 hippocampal cultures. The basal level of Arc/Arg3.1 protein was low but increased significantly in both the soma and dendrites during the 5 min incubation with DHPG (50  $\mu$ M) (Figure 2A). The increase of Arc/Arg3.1



**Figure 1. Arc/Arg3.1 Is Required for Hippocampal mGluR-LTD**

Field excitatory postsynaptic potentials (fEPSPs) were recorded in the hippocampal Schaffer collateral-CA1 synapses derived from *Arc/Arg3.1* KO mice and compared to WT littermate controls.

(A) Average time course of the change in fEPSP slope induced by the group I mGluR agonist (R,S)-DHPG (50  $\mu$ M, for 5 min). LTD of WT mice was  $72.8\% \pm 2.0\%$  of baseline at  $t = 70$  min ( $n = 10$ ). In *Arc/Arg3.1* KO mice, fEPSPs were  $92.1\% \pm 3.7\%$  of the baseline at  $t = 70$  min ( $n = 9$ ).  $p < 0.001$  when compared to littermate WT. Error bars indicate the standard error of the mean. Measurements correspond to the time points indicated on the time course graph in this and all subsequent figures.

(B) Time course of the change in fEPSP slope produced by paired-pulse low-frequency stimulation (PP-LFS: at 1 Hz, 50 ms interstimulus interval, for 15 min) in the presence of the NMDA receptor antagonist D-APV (50  $\mu$ M). LTD of WT mice was  $79.9\% \pm 2.1\%$  of baseline at  $t = 80$  min ( $n = 12$ ). In *Arc/Arg3.1* KO mice, fEPSPs were  $94.3\% \pm 2.1\%$  of the baseline at  $t = 80$  min ( $n = 13$ ).  $p < 0.0001$ . Scale bars, 0.5 mV/10 ms.

(C) Five minutes of DHPG application resulted in a loss of surface GluR1 at 15 min ( $n = 20$ ,  $***p < 0.005$ ) and 60 min ( $n = 19$ ,  $*p < 0.05$ ) after DHPG application, compared to untreated controls in WT hippocampal cultures. *Arc/Arg3.1* KO neurons did not exhibit any changes in surface GluR1 levels after DHPG treatment. Representative pictures of cultures are shown using an LUT scale where white is high intensity and dark red is low intensity. CTL, control.


(D) Five minutes of DHPG application resulted in an increase of internalized GluR1 at 15 min ( $n = 20$ ,  $*p < 0.05$ ) compared with untreated cultures. *Arc/Arg3.1* KO neurons did not exhibit changes in internalized GluR1 levels after DHPG treatment. Error bars indicate SEM in this and all subsequent figures.

protein was blocked by the protein synthesis inhibitor emetine, indicating a role for de novo translation. The induced Arc/Arg3.1 immunoreactivity in both proximal and distal dendrites was detected within 5 min of mGluR activation, and there was no evidence of a concentration gradient that might occur with rapid transport of Arc/Arg3.1 from the soma. The rapidity and distribution of the response suggests that Arc/Arg3.1 is synthesized locally in dendrites and is consistent with the observation that mGluR-LTD is expressed in isolated dendrites (Huber et al., 2000). Similar levels of Arc/Arg3.1 induction during 5 min incubation of DHPG were observed by western blot analysis using forebrain cultures (Figure S2A). Treatment with BDNF (10 ng/ml) also increased Arc/Arg3.1 protein expression, but, in contrast to DHPG, this was evident only after 40 min (Figure S2A).

The rapid increase of Arc/Arg3.1 protein could be mediated by an enhanced rate of translation or by a stable level of translation together with reduced degradation. As reported previously (Rao

et al., 2006), the proteasome inhibitor MG132 increased Arc/Arg3.1 protein but did not block the ability of DHPG to further increase Arc/Arg3.1 (Figure S2B). Induction of Arc/Arg3.1 by DHPG at 5 min was blocked by 5 min pretreatment of emetine or cycloheximide (Figures 2A7 and 2B). These data support the notion that Arc/Arg3.1 induction following DHPG treatment involves an increase in the rate of de novo protein translation.

To examine the possible role of de novo transcription of Arc/Arg3.1 mRNA, we monitored the effect of the transcription blocker actinomycin D. Actinomycin D (10  $\mu$ M, 5 min pretreatment and 5 min with or without DHPG) did not alter the DHPG-induced increase of Arc/Arg3.1 protein (Figures 2C and 2E). DHPG did evoke a modest increase of Arc/Arg3.1 mRNA, but this was detected only after 20 min (Figure 2F). The time course of the delayed Arc/Arg3.1 protein expression by DHPG or BDNF correlated with the mRNA induction, and actinomycin D blocked this response (data not shown). These observations



**Figure 2. Arc/Arg3.1 Protein Is Rapidly Synthesized by Group I mGluR Activation**

(A) Stimulation of hippocampal neurons with DHPG (50 μM) for 5 min increased Arc/Arg3.1 immunoreactivity in both cell body ( $1.34 \pm 0.063$  of untreated soma,  $n = 13$ ) and dendrites ( $1.58 \pm 0.095$  of untreated dendrites,  $n = 38$ ). The rapid increase of Arc/Arg3.1 was blocked by the protein synthesis inhibitor emetine (10 ng/ml, 10 min).

(B) High-dose cycloheximide (CHX, 50 μM, total 10 min: 5 min pretreatment and 5 min with or without DHPG) blocked the induction of Arc/Arg3.1 by DHPG (5 min).

(C) Transcription inhibitor actinomycin D (ActD: 10 μM, 5 min pretreatment and 5 min with or without DHPG) did not block the induction of Arc/Arg3.1 by DHPG (5 min).

(D) Low-dose CHX increased the level of Arc/Arg3.1 protein. Neurons were treated with vehicle or various doses of CHX for 10 min. Total protein synthesis was measured by counting the incorporation of <sup>35</sup>S methionine and cysteine in TCA precipitant.

(E) Statistical analysis of western blots. Five minute treatment of DHPG significantly increased the level of Arc/Arg3.1. Inhibition of new protein synthesis by high dose of cycloheximide not only blocked the induction of Arc/Arg3.1 protein but also slightly decreased the level of Arc/Arg3.1 upon stimulation with DHPG. Inhibition of transcription by actinomycin D did not affect the level of Arc/Arg3.1. Low-dose CHX (50–100 nM, 5 min pretreatment and 5 min with or without DHPG) increased the level of Arc/Arg3.1, which was not further induced by DHPG. \* $p < 0.05$ , \*\* $p < 0.01$ .

(F) The level of Arc/Arg3.1 mRNA was measured using real-time RT-PCR. Stimulation of neurons with BDNF (10 ng/ml) and forskolin (50 μM) induced the level of Arc/Arg3.1 mRNA 40 min and 20 min after stimulation, respectively. DHPG slightly increased the level of Arc/Arg3.1 mRNA at 20 and 40 min after stimulation.

\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.005$ .

suggest that the rapid increase in de novo translation requires Arc/Arg3.1 mRNA that is present in neurons prior to DHPG stimulation, while the delayed Arc/Arg3.1 expression is coupled to de novo transcription. We note that Arc/Arg3.1 mRNA is detected in dendrites of unstimulated cultured neurons (Giorgi et al., 2007), and we detect Arc/Arg3.1 mRNA in stratum radiatum of the hippocampal CA1 region from home-caged mice (Figure S3).


### Low Dose Cycloheximide Can Increase Arc/Arg3.1 Protein Expression

In examining the dose dependence of cycloheximide's actions, we noted that the level of Arc/Arg3.1 protein rapidly increased when neurons are treated with low doses (Figure 2D). For example, 100 nM cycloheximide increased Arc/Arg3.1 protein within 10 min. Even at these low doses, cycloheximide effectively reduced general protein synthesis. 100 nM cycloheximide reduced the total incorporation of <sup>35</sup>S-labeled methionine and cysteine into TCA precipitant to ~60%. Previous studies have noted the

paradoxical action of low-dose cycloheximide to increase the synthesis of specific proteins and rationalized this action by suggesting that global reduction of elongation can increase the availability of factors that are required for translation initiation of specific transcripts that are poorly initiated under control conditions (Fernandez et al., 2005; Gupta and Ono, 1997; Perlman and Feldman, 1982; Scheetz et al., 2000; Walden and Thach, 1986). This notion contributed to our analysis of the eEF2 pathway (below), since activated eEF2 inhibits elongation and can paradoxically increase translation of certain mRNAs (Chotiner et al., 2003; Scheetz et al., 2000).

### eEF2K Physically Associates with Homer and Group I mGluRs

Homer proteins bind group I mGluRs and play a role in their signaling by also binding signaling partners, including IP<sub>3</sub>R (Tu et al., 1998). Homer proteins bind two known sequence motifs: PPxxF (type 1) and LPSSPSS (type 2) (Yuan et al., 2003). When we searched for candidate Homer-binding



**Figure 3. eEF2K Binds Homer and mGluR1/5**

(A) Schematic diagram of eEF2K. The N terminus of eEF2K contains a  $\text{Ca}^{2+}$ /calmodulin (CaM) binding motif and an  $\alpha$ -kinase domain. The C-terminal eEF2 targeting domain, which recruits the substrate, eEF2, is linked to the hyperphosphorylated internal region. Putative Homer binding site is shown above the diagram.

(B) Coimmunoprecipitation (co-IP) of eEF2K and Homer. HA-tagged (HA-) eEF2K was coexpressed with myc-tagged WT, W27A, or G91N Homer3 in HEK293T cells, and IP was performed with anti-myc antibody. HA-eEF2K co-IPed with WT or W27A Homer 3 was coexpressed, but not with G91N Homer.

(C) mGluR5 increases the interaction of eEF2K and Homer. HA-eEF2K was transfected with or without HA-mGluR5. IP was performed by anti-Homer2 antibody, which IPed endogenous Homer2 protein. Western blot was performed using anti-HA antibody. Co-IP of HA-eEF2K was increased when mGluR5 was coexpressed.

(D) eEF2K co-IPs with mGluR5. HEK293T cells were transfected with HA-eEF2K with or without HA-mGluR5, and lysates were IPed with anti-mGluR5 antibody and blotted with anti-HA antibody. eEF2K co-IPed only when mGluR was coexpressed. Samples were boiled before loading to aggregate and separate mGluR5 monomer from eEF2K.

(E) mGluR1 co-IPs with eEF2K. HEK293T cells were transfected with mGluR1 and eEF2K, and lysates were IPed with mycAb. Samples were not boiled to show the monomer of mGluRs.


(F and G) mGluR2 and mGluR4 do not co-IP with eEF2K.

molecules (<http://us.expasy.org/cgi-bin/scanprosite>), we found that eEF2K possesses a type 2 Homer-binding motif (Figure 3A). eEF2K is a highly conserved enzyme that phosphorylates and regulates its only known substrate, eEF2 (Ryzanov et al., 1988). The N-terminal half of eEF2K contains a  $\text{Ca}^{2+}$ -calmodulin (CaM) binding site, which is required for its activation, and an  $\alpha$  kinase domain. The C-terminal half of eEF2K functions as a targeting domain that is required for it to phosphorylate eEF2. A linker region between the N and C terminus includes the putative Homer-binding site and is phosphorylated by multiple signaling kinases, including PI3K/mTOR/S6K, MAPK, and PKA (Browne and Proud, 2002).

eEF2K and Homer coimmunoprecipitated (co-IP) from HEK293T cells (Figure 3B). The EVH1 domain of Homer is required to bind eEF2K, and mutation of a critical binding surface for polyproline ligands (Homer3 G91N) disrupted binding. As anticipated by the conservation of their EVH1 domains, Homer 1, -2, and -3 bind eEF2K (data not shown).

Homer. The interaction of eEF2K and group I mGluRs was observed even when Homer was not coexpressed (Figure 3D and 3E), and eEF2K bound to mGluR5 mutants that do not bind Homer (Figures S4A and S4B). eEF2K also co-IPed with mGluR1 (Figure 3E), another member of group I mGluRs, but not with other mGluRs, including mGluR2 and mGluR4 (Figure 3F and 3G).

To identify regions of eEF2K that are critical for binding Homer and mGluR, we performed a deletion analysis of eEF2K (Figures S4C and S4D). The linker region of eEF2K, which includes the type 2 Homer ligand, appears essential for binding Homer, since N-terminal fragments that include this region bind, while C-terminal fragments or N-terminal fragments that do not include the linker region do not bind. eEF2K binding to mGluR5 appears to be more complex, since both N- and C-terminal fragments of eEF2K bind mGluR5 (Figure S4C). These data suggest that mGluR, Homer, and eEF2K assemble by multiple interactions into a tertiary complex.



**Figure 4. Dynamic Interaction of eEF2K and mGluR5**

(A) Calcium dissociates eEF2K from mGluR5. HEK293T cells were transfected with HA-eEF2K with or without myc-mGluR5, and cells were harvested with lysis buffer without calcium or containing various concentrations of free calcium. Calmodulin (CaM) (25 μg/ml) was also added to the lysis buffer as indicated. Binding was decreased at [Ca<sup>2+</sup>] higher than 10 μM.

(B) Phospho-eEF2 was not detected in the hippocampus of *eEF2K* KO, while the level of total eEF2, GluR1, GluR2/3, mGluR5, α-CaMKII, Arc/Arg3.1, and actin was not altered in *eEF2K* KO mice compared to WT littermate controls.

(C) Synaptoneurosomes, prepared from the forebrain of *eEF2K* KO and WT mice, were stimulated with vehicle or DHPG for 20 min. Synaptoneurosomes were then lysed and immunoprecipitated with anti-eEF2K antibody. mGluR5 co-IPed with eEF2K only in WT samples. Stimulation of synaptoneurosomes with DHPG decreased the co-IP of mGluR5.

### The Interaction of eEF2K with mGluR Is Dynamic and Is Modulated by Ca<sup>2+</sup> and mGluR Activity

The kinase activity of eEF2K is known to be regulated by Ca<sup>2+</sup> via its Ca<sup>2+</sup>-CaM-binding domain (Nairn and Palfrey, 1987; Ryazanov, 1987). To test whether Ca<sup>2+</sup> modulates the mGluR5-eEF2K binding, co-IP experiments were performed using lysates from cotransfected HEK293T cells in the presence of defined concentrations of free Ca<sup>2+</sup> (Figure 4A). Co-IP was robust at [Ca<sup>2+</sup>] less than 1 μM but markedly decreased at concentrations >10 μM. mGluR5 binding to a C-terminal fragment of eEF2K that lacks the CaM-binding domain but retains binding to mGluR5 was not inhibited by [Ca<sup>2+</sup>] (Figure S4E). These results indicate that [Ca<sup>2+</sup>] can modulate the interaction of group I mGluRs with eEF2K and suggest a role for CaM binding to eEF2K.

We used *eEF2K* KO mice in our analysis of mGluR-eEF2K binding. *eEF2K* KO mice were viable and fertile (Ryazanov, 2002) and showed the anticipated absence of phosphorylated eEF2 (Thr56) (Figure 4B). The levels of several synaptic proteins were not altered in the hippocampus of KO mice (Figure 4B). Synaptoneurosomes from forebrains of WT and *eEF2K* KO mice were prepared and stimulated with DHPG for 20 min. Co-IP experiments using anti-eEF2K antibody confirmed that native mGluR5 associated with eEF2K (Figure 4C). The co-IP of mGluR5 was reduced when synaptoneurosomes were stimulated with DHPG. Interaction of endogenous mGluR5 and eEF2K was also reduced upon DHPG stimulation of cultured neurons (Figure S4F). We conclude that mGluR and eEF2K associate in vivo and that their interaction is reduced by mGluR activation.

### Group I mGluRs Dynamically Regulate the Phosphorylation of eEF2


Activated eEF2K selectively phosphorylates eEF2 (Ryazanov et al., 1988). To assess whether mGluR activates this pathway in conditions that evoke LTD, we monitored the level of phospho-eEF2 in hippocampal slices of either WT or *eEF2K* KO mice using the same stimulus parameters that induce mGluR-LTD. Activation of mGluR increased the phosphorylation of eEF2 in the stratum pyramidale (s.p.), and stratum radiatum (s.r.) of the hippocampal CA1 region within 5 min (Figure 5A). By 30 min after washout of DHPG, the level of phospho-eEF2 was reduced to prestimulation level. No phosphorylation of eEF2 was detected in *eEF2K* KO slices. The transient induction of phospho-eEF2 by DHPG was confirmed by western blot analysis in hippocampal slices (Figure S5A).

To further examine dendritic localization of eEF2K activity, DIV14 neurons were stimulated with DHPG for 5 min and stained with phospho-eEF2 and PSD95, a marker for excitatory synapses (Figure 5B). Phospho-eEF2 was present in dendritic shafts and the cell body. Phospho-eEF2 also showed a distinct punctal distribution in spines that colocalized with PSD95. Staining was absent in *eEF2K* KO cultures (data not shown). This result is consistent with a previous report that translational regulators, including eEF2K, are enriched in synaptic fractions (Asaki et al., 2003).

Phosphorylation of eEF2 is known to inhibit translational elongation. Therefore, we examined the prediction that global protein translation might be transiently reduced co-incident with the transient increase of phospho-eEF2. Stimulation of neurons with DHPG for 5 min transiently decreased the incorporation of <sup>35</sup>S amino acids into TCA precipitants, and this effect was reversed 20 min after washout of DHPG (Figure S5B). A previous study reported that DHPG rapidly increased protein synthesis in synaptoneurosomes (Weiler et al., 2004). We observed that DHPG did not induce p-eEF2 in synaptoneurosomes (data not shown), and it is possible that the eEF2-dependent translational mechanism is not maintained in broken cell preparations.

### Rapid De Novo Arc/Arg3.1 Translation Is Selectively Absent in *eEF2K* KO Neurons

Arc/Arg3.1 expression was examined in DIV14 forebrain neuronal cultures prepared from WT and *eEF2K* KO mice. The



**Figure 5. Rapid Induction of Arc/Arg3.1 by Group I mGluRs Is Dependent on eEF2K**


(A) Hippocampal slices were prepared from WT and *eEF2K* KO mice and were stimulated with DHPG for 5 min. phospho-eEF2 (p-eEF2, red) in area CA1 was increased by DHPG within 5 min and declined by 30 min following washout. Specificity of phospho-eEF2 was confirmed by staining of *eEF2K* KO slices. s.p., stratum pyramidale; s.r., stratum radiatum.

(B) Cultured hippocampal neurons were treated with DHPG for 5 min and stained with phospho-eEF2 (red) and PSD95 (green) antibodies on DIV14. phospho-eEF2 showed punctal distribution in dendritic spines and dendritic shafts. phospho-eEF2 in spines colocalized with PSD95 (arrows). (B2), (B3), and (B4) are enlarged images of the rectangular region of (B1).

(C and D) mGluR-dependent rapid synthesis of Arc/Arg3.1 is absent in *eEF2K* KO neurons. Neurons from the forebrains of WT or *eEF2K* KO mice were cultured for DIV14 and treated with DHPG (50  $\mu$ M, 5 min). Phosphorylation of eEF2 was undetectable in *eEF2K* KO neurons. No difference in the level of mGluR5 was observed between WT and *eEF2K* KO neurons. An arrowhead indicates a non-specific band. p values were obtained by paired t test comparing basal and drug-treated levels. p values for comparison of WT and *eEF2K* KO mice were obtained by Student's t test. \*p < 0.05, \*\*p < 0.01, n = 8. Error bars are SEM.

(E) Arc/Arg3.1 mRNA expression is not altered in *eEF2K* KO neurons. The level of Arc/Arg3.1 mRNA was measured in WT and *eEF2K* KO neurons following the stimulation with DHPG.


(F) Low-dose cycloheximide (CHX) increases Arc/Arg3.1 protein expression. Cultured *eEF2K* KO neurons were treated with indicated doses of CHX for 10 min. \*p < 0.05, n = 8.



steady-state expression of Arc/Arg3.1 protein was identical in WT and *eEF2K* KO neurons; however, the increase in Arc/Arg3.1 protein 5 min after DHPG in WT neurons was absent in *eEF2K* KO neurons in both biochemical (Figures 5C and 5D) and immunocytochemical assays (Figure 5B). By contrast, Arc/Arg3.1 protein was induced to the same extent in WT and *eEF2K* KO neurons 60 min after DHPG stimulation. Arc/Arg3.1 mRNA was identical in WT and *eEF2K* KO neurons prior to application of DHPG and increased identically at 40 min after stimulation in both WT and *eEF2K* KO neurons (Figure 5E). Accordingly, the lack of rapid induction of Arc/Arg3.1 protein in the *eEF2K* KO neurons is not due to reduced Arc/Arg3.1 mRNA expression. We also note that mGluR signaling that is required for induction of Arc/Arg3.1 mRNA and the delayed increase of Arc/Arg3.1 protein are intact in *eEF2K* KO neurons. Moreover,

If the failure of DHPG to induce rapid synthesis of Arc/Arg3.1 protein in the *eEF2K* KO neurons is due to a selective interruption of the action of phospho-eEF2, we predicted that a low dose of cycloheximide, which does not require eEF2K or phospho-eEF2 to inhibit the elongation step, should induce the synthesis of Arc/Arg3.1 protein in *eEF2K* KO neurons. Treatment of DIV14 *eEF2K* KO neurons with low-dose cycloheximide (50 nM and 100 nM) increased the level of Arc/Arg3.1 protein in *eEF2K* KO neurons (Figure 5F), similar to WT neurons (Figure 2D). High-dose cycloheximide (>1  $\mu$ M) did not induce Arc/Arg3.1 in either WT and *eEF2K* KO neurons. The ability of low-dose cycloheximide to rescue rapid Arc/Arg3.1 induction indicates that mechanisms that mediate rapid Arc/Arg3.1 translation subsequent to inhibition of elongation are intact in *eEF2K* KO neurons.

Arc/Arg3.1 protein expression is identical in the hippocampus of WT and *eEF2K* KO mice (Figure 4B), indicating that eEF2K is not required for basal expression of Arc/Arg3.1 protein in vivo.



**Figure 6. mGluR-LTD Is Impaired in Hippocampal Slices Derived from eEF2K KO Mice**

fEPSPs were recorded in the hippocampal CA1 region of slices derived from eEF2K KO mice and compared to WT littermate controls.

(A) Time course of the change in fEPSP slope produced by paired-pulse low-frequency stimulation (PP-LFS: at 1 Hz, 50 ms interstimulus interval, for 15 min) in the presence of D-APV (50  $\mu$ M). LTD of WT mice was  $77.0\% \pm 2.1\%$  of baseline at  $t = 75$  min ( $n = 13$ ). In eEF2K KO mice, fEPSPs were  $97.5\% \pm 2.4\%$  of baseline  $t = 75$  min ( $n = 15$ ) ( $p < 0.0001$ ).

(B) Time course of the change in fEPSP slope by low-frequency stimulation (LFS: 1 Hz for 15 min). This form of NMDAR-dependent LTD was not altered in eEF2K KO hippocampal slices ( $72.7\% \pm 2.2\%$  of baseline at  $t = 75$  min,  $n = 9$ ) compared to WT ( $73.1\% \pm 3.4\%$  of baseline at  $t = 75$  min,  $n = 7$ ) ( $p > 0.5$ ).

(C) Late-phase of LTP was induced by four stimulus trains (100 Hz each) with an intertrain interval of 3 s. In WT, fEPSPs were increased to  $171.5\% \pm 13.4\%$  of baseline immediately after stimulation ( $t = 30$  min) and were sustained at the level of  $138.4\% \pm 7.7\%$  of baseline at  $t = 175$  min ( $n = 6$ ). However, in eEF2K KO, the initial LTP ( $204.6\% \pm 8.9\%$  of baseline at  $t = 30$  min) was maintained for 3 hr after stimulation ( $200.1\% \pm 11.9\%$  of baseline at  $t = 175$  min,  $n = 5$ ). LTP was significantly greater in slices derived from eEF2K KO mice compared to those from WT mice at this time point ( $p < 0.005$ ).

(D) Average time course of the change in fEPSP slope induced by DHPG (50  $\mu$ M, for 5 min). LTD of WT mice was  $64.7\% \pm 5.2\%$  of baseline at  $t = 90$  min ( $n = 7$ ). In eEF2K KO mice, LTD was significantly impaired ( $108.7\% \pm 3.6\%$  of baseline at  $t = 90$  min,  $n = 8$ ). Treatment with low-dose cycloheximide (LD-CHX, 50–75 nM) for 10 min starting from 5 min prior to DHPG restored DHPG-LTD in eEF2K KO ( $75.7\% \pm 7.4\%$ ,  $n = 5$ ). In WT mice, treatment with LD-CHX did not alter the expression of LTD ( $69.0\% \pm 2.6\%$ ,  $n = 5$ ).  $p < 0.001$  when eEF2K KO DHPG only was compared to eEF2K KO LD-CHX + DHPG, WT DHPG only, or WT LD-CHX + DHPG. Scale bars, 0.5 mV/10 ms.

### mGluR-LTD and PP-LFS LTD Are Selectively Absent in eEF2K KO Hippocampal Slices

The role of eEF2K in plasticity of the Schaffer collateral-CA1 synapse was tested using acute hippocampal slices. Baseline measures of synaptic strength and presynaptic function were not altered in the eEF2K KO slices (Figure S7). However, LTD induced by either PP-LFS ( $97.5\% \pm 2.4\%$  of baseline) or DHPG ( $108.7\% \pm 3.6\%$  of baseline) was impaired in the eEF2K KO slices (Figures 6A and 6D). The immediate short-term synaptic depression following DHPG stimulation was identical in WT and eEF2K KO slices; however, synaptic strength returned to near baseline levels in the eEF2K KO slices. Similarly, synaptic transmission returned to near baseline levels within 10 min of completion of the PP-LFS protocol. In contrast to the marked deficit of mGluR-dependent LTD, NMDAR-dependent LTD was identical in time course and stability in slices derived from

eEF2K KO mice ( $72.7\% \pm 2.2\%$  of baseline) compared to WT mice ( $73.1\% \pm 3.4\%$  of baseline) (Figure 6B). LTP was also preserved (Figure 6C). LTP of Schaffer collateral-CA1 synapses was induced by four trains of high-frequency stimulation with an intertrain interval of 3 s. In WT slices, fEPSP was increased to  $171.5\% \pm 13.4\%$  of baseline immediately after stimulation ( $t = 30$  min) and sustained at the level of  $138.4\% \pm 7.7\%$  of baseline at  $t = 175$  min. These stimulus parameters are reported to evoke a form of synaptic plasticity that requires de novo protein synthesis for maintenance longer than  $\sim 60$  min and is referred to as late LTP (L-LTP) (Kelleher et al., 2004). In slices prepared from eEF2K KO mice, the initial induction was  $204.6\% \pm 8.9\%$  of baseline at  $t = 30$  min, and this was sustained for 3 hr after stimulation ( $200.1\% \pm 11.9\%$  of baseline at  $t = 175$  min) (Figure 6C). The magnitude of LTP was significantly greater in eEF2K KO than WT after 30 min of induction ( $p < 0.005$ ). These results indicate



that *eEF2K* KO disrupts mGluR-LTD but does not alter NMDAR-dependent LTD or early LTP. The apparent enhancement of late-phase LTP deserves further study.

Our proposed mechanism for the mGluR-LTD deficit in the *eEF2K* KO slices is linked to failure to rapidly translate Arc/Arg3.1. Since low-dose cycloheximide induced Arc/Arg3.1 synthesis and did not depend on phospho-eEF2, we examined the possibility that cycloheximide could rescue mGluR-LTD in slices from *eEF2K* KO mice. We found that a 10 min exposure to 50–75 nM cycloheximide (low-dose CHX: LD-CHX) beginning 5 min prior to addition of DHPG rescued mGluR-dependent LTD in the *eEF2K* KO slice ( $75.7\% \pm 7.4\%$  of baseline,  $p < 0.001$ , compared to DHPG only in *eEF2K* KO slices) (Figure 6D). The same treatment of WT slices did not substantially alter the time course of mGluR-LTD ( $69.0\% \pm 2.6\%$  of baseline,  $p > 0.5$ , compared to DHPG only in WT slices). Low-dose cycloheximide had no effect on baseline synaptic transmission in the absence of mGluR stimulation ( $101.2\% \pm 2.0\%$  for WT slices;  $100.4\% \pm 4.6\%$  for *eEF2K* KO slices). These observations confirm that mGluR signaling required for mGluR-LTD is selectively impaired in *eEF2K* KO in a manner that can be rescued by transient application of low-dose cycloheximide.

#### **mGluR-LTD, but Not Homeostatic Plasticity, Is Disrupted in *eEF2K* KO Neurons in Culture**

To further assess the selectivity of the *eEF2K* KO effect on neuronal function, we examined two forms of neuronal plasticity that can be assayed in primary neuronal cultures. Treatment of cultures with DHPG for 5 min to evoke mGluR-LTD reduced the ratio of surface to total GluR2/3 by  $\sim 30\%$  in WT neurons but did not significantly reduce this measure in *eEF2K* KO neurons (Figures S8A and S8B). This result parallels the deficit of mGluR-LTD seen in acute slices. Cultures were also assayed for homeostatic adaptations of surface AMPA receptors since this response is markedly altered in *Arc/Arg3.1* KO neurons (Shepherd et al., 2006). Treatment of *eEF2K* KO cortical cultures for 2 days with either tetrodotoxin (TTX) or bicuculline evoked homeostatic increases and decreases of surface GluR1 that were identical to WT neurons (Figure S8C). Thus, *eEF2K* KO results in a selective disruption of mGluR-dependent LTD.


#### ***Fmr1* KO Disrupts Rapid, but Not Delayed, Induction of Arc/Arg3.1 Protein**

We examined the role of the eEF2K/eEF2/Arc mechanism in the aberrant plasticity described in *Fmr1* KO mice. FMRP has been reported to bind Arc/Arg3.1 mRNA (Iacoangeli et al., 2008; Zalfa et al., 2003) and is hypothesized to inhibit translation prior to mGluR stimulation (Bear et al., 2004; Narayanan et al., 2007). To assess whether FMRP might be critical for either rapid or delayed induction of Arc/Arg3.1 protein following mGluR stimulation, we prepared primary neuronal cultures from *Fmr1* KO mice and stimulated with DHPG. Arc/Arg3.1 expression in unstimulated cultures was not consistently different between WT and *Fmr1* KO neurons. Moreover, Arc/Arg3.1 protein increased 60 min after DHPG stimulation in *Fmr1* KO neurons identically as in WT neurons (Figure 7A). However, the rapid increase of Arc/Arg3.1 protein following DHPG stimulation was absent in *Fmr1* KO neurons (Figure 7A). DHPG activated mGluR/eEF2K

signaling in the *Fmr1* KO neurons since phospho-eEF2 was identically induced as in WT neurons (Figure 7A). Assays of Arc/Arg3.1 protein stability and induction following proteasome inhibition with MG132 did not reveal differences between WT and *Fmr1* KO neurons (Figures S9B and S9C). Biochemical experiments to monitor Arc/Arg3.1 expression using acute hippocampal slices revealed that basal Arc/Arg3.1 expression was highly variable even when normalized to total protein or actin, indicating a limitation of this preparation (Taubenfeld et al., 2002). When examined histochemically, basal Arc/Arg3.1 varied through the thickness of the slice (data not shown). We conclude that *Fmr1* KO neurons selectively lack the ability to rapidly upregulate Arc/Arg3.1 expression. The reported increase of Arc/Arg3.1 mRNA in polysome fractions from *Fmr1* KO mice (Zalfa et al., 2003) suggests that failure to detect a DHPG-evoked rapid increase of Arc/Arg3.1 protein is linked to elevated constitutive expression.

#### **Arc/Arg3.1 Is Required for mGluR-LTD and PP-LFS LTD in *Fmr1* KO Mice**

In anticipation of physiological studies to assess the role of Arc/Arg3.1 in synaptic plasticity of *Fmr1* KO mice, we examined Arc/Arg3.1 protein expression in hippocampus. Arc/Arg3.1 protein has previously been reported to be modestly upregulated in both total brain and synaptosomal fractions of *Fmr1* KO mice (Zalfa et al., 2003). However, in our assays, Arc/Arg3.1 protein was not consistently different in hippocampus (either in vivo or in acute slices) or cortex when care was taken to sacrifice mice without behavioral activation. We generated mice in which both *Fmr1* (in FVB background) and *Arc/Arg3.1* (in B6 background) were deleted. Double *Arc/Arg3.1/Fmr1* KO (DKO) mice are viable, fertile, and not different from WT mice in size or postnatal survival. Indices of basal synaptic transmission were normal in *Fmr1* KO and *Arc/Arg3.1/Fmr1* DKO (Figures S9D and S9E). As reported previously (Nosyreva and Huber, 2006), DHPG evoked a sustained reduction of synaptic strength ( $68.2\% \pm 2.6\%$  of baseline for *Fmr1* single KO slices;  $73.0\% \pm 6.6\%$  of baseline for FVB WT slices; Figure 7C). Jackson laboratory provides *Fmr1* KO mice in the FVB background, and the magnitude of LTD was not significantly different from FVB WT mice. As reported previously in studies of *Fmr1* KO in the B6 background, mGluR-LTD was not inhibited by high-dose cycloheximide (60  $\mu$ M) (Figure 7B). In *Arc/Arg3.1/Fmr1* DKO (in B6/FVB), DHPG evoked an initial reduction of synaptic strength that was not different from WT, *Arc/Arg3.1* KO, or *Fmr1* KO. However, expression of DHPG-evoked LTD was significantly impaired in *Arc/Arg3.1/Fmr1* DKO ( $85.9\% \pm 4.1\%$ ,  $p < 0.01$  compared to *Fmr1* single KO or FVB WT). PP-LFS LTD was also impaired in *Arc/Arg3.1/Fmr1* DKO ( $88.3\% \pm 2.1\%$  of baseline for *Arc/Arg3.1/Fmr1* DKO slices;  $75.5\% \pm 3.7\%$  of baseline for *Fmr1* single KO;  $80.5\% \pm 2.6\%$  of baseline for FVB WT slices,  $p < 0.05$  when *Arc/Arg3.1/Fmr1* DKO was compared to *Fmr1* single KO or FVB WT; Figure 7D). These results indicate that Arc/Arg3.1 is required for mGluR-LTD in both WT and *Fmr1* KO neurons. Deletion of Arc/Arg3.1 does not entirely prevent DHPG or PP-LFS LTD, suggesting that additional mechanisms contribute to the aberrant LTD in *Fmr1* KO mice.



**Figure 7. LTD Is Impaired in Hippocampal Slices Derived from *Arc/Arg3.1/Fmr1* Double KO Mice**

(A) DIV14 *Fmr1* KO neurons were treated with DHPG as indicated in Figure 5C. Rapid but not delayed synthesis of *Arc/Arg3.1* was absent in *Fmr1* KO. The regulation of phospho-eEF2 was intact in *Fmr1* KO neurons.

(B) High-dose cycloheximide (60  $\mu$ M; HD-CHX) did not block DHPG-LTD of *Fmr1* KO slices. In the presence of a high dose of cycloheximide, DHPG-LTD of *Fmr1* KO was  $72.3\% \pm 4.8\%$  of baseline at  $t = 105$  min ( $n = 5$ ), while DHPG-LTD in WT (FVB) slices was blocked (fEPSP was  $95.5\% \pm 2.9\%$  of baseline at  $t = 105$  min ( $n = 4$ );  $p < 0.01$  when *Fmr1* KO was compared to FVB WT).

(C) Average time course of fEPSP slope of *Arc/Arg3.1/Fmr1* double KO (DKO) mice. mGluR-LTD was induced by DHPG (50  $\mu$ M, for 5 min). DHPG-LTD of *Arc/Arg3.1/Fmr1* DKO was  $85.9\% \pm 4.1\%$  of baseline at  $t = 75$  min ( $n = 8$ ). In *Fmr1* KO, DHPG-LTD was  $68.2\% \pm 2.6\%$  of baseline at  $t = 75$  min ( $n = 6$ ). In WT, DHPG-LTD was  $73.0\% \pm 6.6\%$  of baseline at  $t = 75$  min ( $n = 5$ ).  $p < 0.01$  when *Arc/Arg3.1/Fmr1* DKO was compared to either WT or *Fmr1* KO. fEPSPs of post-DHPG in *Fmr1* KO were not significantly different from those in FVB WT.

(D) Time course of the change in fEPSP slope by PP-LFS. PP-LFS LTD of *Arc/Arg3.1/Fmr1* DKO was  $88.3\% \pm 2.1\%$  of baseline at  $t = 65$  min ( $n = 6$ ). In *Fmr1* KO, PP-LFS LTD was  $75.5\% \pm 3.7\%$  of baseline at  $t = 65$  min ( $n = 8$ ). In FVB WT, PP-LFS LTD was  $80.5\% \pm 2.6\%$  of baseline at  $t = 65$  min ( $n = 8$ ).  $p < 0.05$  when *Arc/Arg3.1/Fmr1* DKO was compared to either WT or *Fmr1* KO. fEPSPs of post-DHPG in *Fmr1* KO were not significantly different from those in FVB WT ( $p = 0.4$ ).

Scale bars, 0.5 mV/10 ms.

## DISCUSSION


### *Arc/Arg3.1* Plays a Central Role in mGluR-Dependent LTD

The present study identifies *Arc/Arg3.1* as a molecule that is required for certain forms of LTD that are known to be dependent on rapid, de novo protein synthesis, including mGluR-LTD and PP-LFS-LTD. Upon activation of group I mGluRs, *Arc/Arg3.1* protein is rapidly upregulated in a process that requires de novo translation. *Arc/Arg3.1* protein functions as a regulatory factor for a postsynaptic endocytic signaling pathway that includes endophilins 2 and -3 and dynamin, which together mediate selective endocytosis of AMPA type glutamate receptors (Chowdhury et al., 2006). Accordingly, the local and rapid increase of *Arc/Arg3.1* protein can be linked to the selective downregulation of AMPAR at specific synapses during LTD (Figure 8).

While group I mGluR activation results in a modest rise of *Arc/Arg3.1* mRNA, the translation that underlies mGluR-LTD

utilizes *Arc/Arg3.1* mRNA that is present in neurons prior to stimulation. *Arc/Arg3.1* is regulated as an immediate-early gene that is strongly induced during learning-related behaviors (Guzowski et al., 1999) and expressed in discrete populations of neurons that are part of behaviorally activated networks even when rodents are resting in home cage (Marrone et al., 2008). *Arc/Arg3.1* mRNA is present in dendritic regions of CA1 neurons of mice sacrificed immediately from their home cage (Figure S3), indicating that basal *Arc/Arg3.1* mRNA can be either activity independent or related to prior neuronal activity. The rapid translation of dendritic *Arc/Arg3.1* mRNA rationalizes the known dependence of mGluR-LTD on rapid de novo protein synthesis and independence of de novo RNA synthesis (Huber et al., 2000). It also suggests that prior neuronal activity that drives the expression of *Arc/Arg3.1* mRNA could modify subsequent mGluR-dependent plasticity.

In further support for the notion that *Arc/Arg3.1* is an essential LTD molecule, we find that mGluR-LTD in *Fmr1* KO mice is also



**Figure 8. eEF2K, FMRP, and Rapid De Novo Translation of Arc/Arg3.1 Protein in mGluR-LTD**

Group I mGluRs activate eEF2K via calcium-calmodulin (CaM). eEF2K phosphorylates eEF2, which inhibits elongation generally but increases Arc/Arg3.1 translation. Arc/Arg3.1 forms a complex with endophilin2/3 (Endo) and dynamin (Dyn) and induces the internalization of AMPAR (Chowdhury et al., 2006). FMRP inhibits the translation of Arc/Arg3.1 at the basal state. Arc/Arg3.1 induction alone is not sufficient for mGluR-LTD, indicating that mGluR activates another pathway that is required to internalize AMPAR (Cho et al., 2008). In *Fmr1* KO mice, the synthesis of Arc/Arg3.1 protein is constitutively derepressed, and de novo synthesis of Arc/Arg3.1 is not required for mGluR-LTD.

all mRNAs and appears central to its role in regulating Arc/Arg3.1 protein translation since low-dose cycloheximide, which mimics phospho-eEF2 by inhibiting translation elongation, can also induce Arc/Arg3.1 expression. Low-dose cycloheximide has been reported to increase the translation of mRNAs that are poorly initiated in basal conditions, including  $\alpha$ -CaMKII (Fernandez et al., 2005; Gupta and Ono, 1997; Perlman and Feldman, 1982; Scheetz et al., 2000; Walden and Thach, 1986). Both  $\alpha$ -CaMKII and Arc/Arg3.1 mRNAs are localized to dendrites (Mori et al., 2000;

Steward et al., 1998) and contain internal ribosome entry site (IRES) sequences in their 5' untranslated regions (Pinkstaff et al., 2001), which may allow the initiation of translation by both Cap-dependent and Cap-independent mechanisms (Pestova et al., 2001). Cap-dependent initiation of translation is tightly controlled by initiation factors such as eukaryotic initiation factor 4E (eIF4E) and eIF4E-binding protein (4EBP). By contrast, Cap-independent initiation through IRES sequence may allow synthesis of proteins when cap-dependent initiation factors are not fully active (Pestova et al., 2001). The seemingly paradoxical effect of phospho-eEF2 and cycloheximide to increase Arc/Arg3.1 protein translation is consistent with a model in which inhibition of global elongation results in a rapid increase in factors that are rate limiting for translation initiation of mRNAs that may be inefficiently initiated under basal conditions (Fernandez et al., 2005; Scheetz et al., 2000; Walden and Thach, 1986). Whether the putative IRES sequence of Arc/Arg3.1 mRNA is required for this response remains to be determined.

**Group I mGluRs and Rapid De Novo Translation; Convergence of eEF2K/eEF2 and FMRP Pathways**

mGluR activation induces the rapid translation of Arc/Arg3.1 and requires eEF2K. eEF2K binds Homer and group I mGluRs and dissociates from this complex in a Ca<sup>2+</sup>-dependent manner. Phospho-eEF2 is present at excitatory synapses and dendrites and is rapidly upregulated in cultured neurons and hippocampal slices by stimulation of group I mGluRs with DHPG. The ability of phospho-eEF2 to inhibit translation is thought to be general for

mGluR activity also rapidly regulates FMRP-dependent signaling in a manner that can enhance rapid, de novo translation (Muddashetty et al., 2007; Narayanan et al., 2007; Westmark and Malter, 2007). mGluR activation results in dephosphorylation of FMRP, and this reduces its inhibitory action on translation (Narayanan et al., 2007). Dephosphorylation of FMRP occurs

within 1 min of mGluR activation and parallels the time course of mGluR-dependent phosphorylation of eEF2. Polysome profiles from WT and *Fmr1* KO brain indicate that FMRP is associated with mRNAs that are not in active translation. In the absence of FMRP, specific mRNAs including *Arc/Arg3.1* shift to fractions that include actively translating polyribosomes (Zalfa et al., 2003). Group I mGluRs are reported to rapidly reduce the histochemical colocalization of FMRP and *Fmr1* mRNA at synapses in association with increased translation of de novo FMRP (Antar et al., 2004). While this is an indirect assay of translational regulation, it supports a model in which mGluR activity reduces FMRP binding to specific mRNAs and thereby provides a dynamic increase in mRNAs in close proximity to synapses for translation. FMRP is not required for mGluR-dependent induction of phospho-eEF2 (Figure 7A), suggesting that mGluR-FMRP and mGluR-eEF2K are parallel pathways. Accordingly, the rapid, de novo *Arc/Arg3.1* translational response required for normal mGluR-LTD appears to require both a transient reversal of FMRP's repressive effect on translation and the general inhibition of the elongation step by phospho-eEF2 (Figure 8).

#### De Novo Translated *Arc/Arg3.1* Is Specifically Required for Normal mGluR-LTD

*Arc/Arg3.1* protein that mediates mGluR-LTD appears to be functionally distinct from the total pool of *Arc/Arg3.1* protein expressed in neurons. For example, the basal expression of *Arc/Arg3.1* protein in *eEF2K* KO neurons is identical to that in WT neurons, yet mGluR-LTD and PP-LFS LTD, which are dependent on *Arc/Arg3.1*, are absent in the *eEF2K* KO. *Arc/Arg3.1* protein in *eEF2K* KO neurons is functional, since homeostatic scaling, which is also dependent on *Arc/Arg3.1* (Shepherd et al., 2006), is intact. This apparent paradox is resolved with the observations that *eEF2K* KO neurons selectively lack rapid, de novo *Arc/Arg3.1* induction, and this deficit is rescued by brief application of low-dose cycloheximide that also rescues mGluR-LTD. Thus, the eEF2-dependent pool of *Arc/Arg3.1* protein appears uniquely capable of mediating LTD. In another example demonstrating the distinct properties of *Arc/Arg3.1* related to its mode of translation, *Arc/Arg3.1* protein is expressed at near normal levels in *Fmr1* KO mice, and *Arc/Arg3.1* protein is induced after 40 min of DHPG stimulation; however, the rapid increase of *Arc/Arg3.1* at 5 min is absent. In this regard, expression of *Arc/Arg3.1* protein in the *Fmr1* KO appears identical to the *eEF2K* KO, but with the technical caveat that we cannot detect changes in basal translation of *Arc/Arg3.1* that are predicted from other studies to be increased in *Fmr1* KO neurons (Zalfa et al., 2003). Our data suggest that an elevation of constitutive *Arc/Arg3.1* translation in *Fmr1* KO underlies mGluR LTD that is aberrantly independent of de novo translation but dependent on *Arc/Arg3.1*. This model rationalizes the distinctly different mGluR-LTD phenotypes in *eEF2K* and *Fmr1* KOs even though they show a similar absence of acute mGluR *Arc/Arg3.1* induction. The relative amount of this eEF2K/FMRP-dependent pool of *Arc/Arg3.1* protein is typically small compared to the *Arc/Arg3.1* level of the delayed response to DHPG or BDNF in neuronal cultures, and we could not detect differences in basal and induced *Arc/Arg3.1* in WT and *eEF2K* KO neurons using conventional histochemical techniques. We infer that proteins

translated by the eEF2K/FMRP-dependent mechanism are uniquely available for mGluR-LTD and anticipate that new imaging methods with high spatial and temporal resolution may shed light on this prediction.

Our studies also indicate that mGluR-LTD requires a second signal in addition to rapid, de novo *Arc/Arg3.1* protein. Experiments that use cycloheximide to rescue mGluR-LTD in the *eEF2K* KO demonstrate that cycloheximide alone is sufficient to induce *Arc/Arg3.1* protein but does not induce LTD unless accompanied by mGluR activation. Similarly, low-dose cycloheximide induces *Arc/Arg3.1* in WT neurons but does not induce LTD without concurrent mGluR activation. These observations rationalize why misregulated expression of *Arc/Arg3.1* protein in the *Fmr1* KO, and presumably other proteins, does not occlude mGluR-LTD. The dual dependence of *Arc/Arg3.1* protein and mGluR activity for aberrant LTD in *Fmr1* KO mice provides a supporting rationale for treatment regimens of Fragile X syndrome that include group I mGluR antagonists (Bear et al., 2004; McBride et al., 2005). In other studies, we have reported that mGluR-LTD requires the rapid cleavage of the neuronal pentraxin receptor protein (NPR) in a process that involves the extracellular metalloprotease tumor necrosis factor- $\alpha$  converting enzyme (TACE/ADAM17) (Cho et al., 2008). The pentraxin domain, released of its transmembrane tether, appears to capture AMPA receptors at the site of endocytosis, and this mechanism is required for mGluR-LTD in both the hippocampus and cerebellum. It is possible that this NPR pathway functions in conjunction with the *Arc/Arg3.1* endosomal pathway and offers a potential new target for agents to modify mGluR-LTD.

#### EXPERIMENTAL PROCEDURES

Constructs, cell culture, real-time RT-PCR, metabolic labeling, and fluorescence in situ hybridization assay are included in Supplemental Experimental Procedures.

#### Antibodies

Anti-phospho-eEF2 (Thr56: rabbit polyclonal) and total-eEF2 (rabbit polyclonal) from Cell Signaling; eEF2K (rabbit polyclonal) and mGluR1 (mouse monoclonal) from BD Biosciences; mGluR5, mGluR2, and PSD-95 from Upstate; mGluR4 from Zymed; horse radish peroxidase (HRP) conjugated HA antibody, HRP-conjugated myc antibody, myc (mouse monoclonal), and actin (mouse monoclonal) from Santa Cruz; *Arc/Arg3.1* (Lyford et al., 1995); N-GluR1 antibody (Shepherd et al., 2006).

#### AMPA Receptor Trafficking Experiments and Immunostaining

Labeling of surface or internalized pool of AMPA receptor was performed as described with minor modifications (Shepherd et al., 2006). Briefly, surface GluR1-containing AMPA receptors were labeled by adding 2.5  $\mu$ g of GluR1-N JH1816 pAb to the neuronal growth media and subsequently incubated at 37°C for 15 or 60 min after 5 min DHPG application. To visualize surface and internalized GluR1, Alexa 555 secondary was added in excess live at 10°C. Neurons were fixed, permeabilized, and subsequently exposed to Alexa 488 secondary to stain internalized receptors (background in the nonpermeabilized control was negligible). Immunocytochemistry of cultured neurons was performed as described (Shepherd et al., 2006). Immunohistochemistry of phospho-eEF2 in WT and *eEF2K* mice was performed as described (Ramirez-Amaya et al., 2005) with slight modifications. See Supplemental Experimental Procedures for details.

**Electrophysiology**

Field recording of excitatory postsynaptic potential (fEPSP) of hippocampal CA1 neurons of postnatal day (P)21–30 male mice as described with minor modifications (Huber et al., 2000). mGluR-LTD was induced by an mGluR1/5 agonist, (R,S)-3,5-DHPG, for 5 min (Tocris, 50  $\mu$ M, unless otherwise indicated), or by paired-pulse low-frequency stimulation (PP-LFS: 50 ms interstimulus interval, 1 Hz, for 15 min) in the presence of D-APV (Tocris, 50  $\mu$ M). NMDAR-dependent LTD was induced by using 900 single pulses delivered at 1 Hz (Huber et al., 2000).

LTP was measured in Schaffer collateral-CA1 synapses in hippocampal slices derived from 8- to 10-week-old male mice. Late-phase LTP (L-LTP) was induced by four trains of high-frequency stimulation (HFS) (100 Hz, 1 s) with 3 s of intertrain interval. See [Supplemental Experimental Procedures](#) for details.

**Western Blotting, Immunoprecipitation Assay, and Surface Biotinylation Assay**

Western blotting, IP assay, and surface biotinylation assay were performed as previously described (Chowdhury et al., 2006; Cho et al., 2008). See [Supplemental Experimental Procedures](#) for details.

**SUPPLEMENTAL DATA**

The Supplemental Data include Supplemental Experimental Procedures and figures and can be found with this article online at <http://www.neuron.org/cgi/content/full/59/1/70/DC1/>.

**ACKNOWLEDGMENTS**

This work was supported by NIMH grants MH053608 (P.F.W.), MH068830 (R.L.H.), MH51106 (D.J.L.), NIDA grant DA00266 (P.F.W.), NIA grant AG019890 (A.G.R.), GM057300 (A.G.R.), and FRAXA (W.K.). We thank Marlin Dehoff and Glory Harris for animal husbandry and David Lieberman for helpful discussions.

Received: August 21, 2007  
Revised: December 22, 2007  
Accepted: May 17, 2008  
Published: July 9, 2008

**REFERENCES**

Antar, L.N., Afroz, R., Dichtenberg, J.B., Carroll, R.C., and Bassell, G.J. (2004). Metabotropic glutamate receptor activation regulates fragile X mental retardation protein and FMR1 mRNA localization differentially in dendrites and at synapses. *J. Neurosci.* *24*, 2648–2655.

Asaki, C., Usuda, N., Nakazawa, A., Kametani, K., and Suzuki, T. (2003). Localization of translational components at the ultramicroscopic level at postsynaptic sites of the rat brain. *Brain Res.* *972*, 168–176.

Bear, M.F., Huber, K.M., and Warren, S.T. (2004). The mGluR theory of fragile X mental retardation. *Trends Neurosci.* *27*, 370–377.

Begueret, J., Perrot, M., and Crouzet, M. (1977). Ribosomal proteins in the fungus *Podospora anserina*: evidence for an electrophoretically altered 60S protein in a cycloheximide resistant mutant. *Mol. Gen. Genet.* *156*, 141–144.

Browne, G.J., and Proud, C.G. (2002). Regulation of peptide-chain elongation in mammalian cells. *Eur. J. Biochem.* *269*, 5360–5368.

Caudy, A.A., Myers, M., Hannon, G.J., and Hammond, S.M. (2002). Fragile X-related protein and VIG associate with the RNA interference machinery. *Genes Dev.* *16*, 2491–2496.

Ceman, S., O'Donnell, W.T., Reed, M., Patton, S., Pohl, J., and Warren, S.T. (2003). Phosphorylation influences the translation state of FMRP-associated polyribosomes. *Hum. Mol. Genet.* *12*, 3295–3305.

Cho, R.W., Park, J.M., Wolff, S.B.E., Xu, D., Hopf, C., Kim, J.-a., Reddy, R.C., Petralia, R.S., Perin, M.S., Linden, D.J., and Worley, P.F. (2008). mGluR1/5-

dependent long-term depression requires the regulated ectodomain cleavage of neuronal pentraxin NPR by TACE. *Neuron* *57*, 858–871.

Chotiner, J.K., Khorasani, H., Nairn, A.C., O'Dell, T.J., and Watson, J.B. (2003). Adenylyl cyclase-dependent form of chemical long-term potentiation triggers translational regulation at the elongation step. *Neuroscience* *116*, 743–752.

Chowdhury, S., Shepherd, J.D., Okuno, H., Lyford, G., Petralia, R.S., Plath, N., Kuhl, D., Huganir, R.L., and Worley, P.F. (2006). Arc/Arg3.1 interacts with the endocytic machinery to regulate AMPA receptor trafficking. *Neuron* *52*, 445–459.

Darnell, J.C., Jensen, K.B., Jin, P., Brown, V., Warren, S.T., and Darnell, R.B. (2001). Fragile X mental retardation protein targets G quartet mRNAs important for neuronal function. *Cell* *107*, 489–499.

Davidkova, G., and Carroll, R.C. (2007). Characterization of the role of microtubule-associated protein 1B in metabotropic glutamate receptor-mediated endocytosis of AMPA receptors in hippocampus. *J. Neurosci.* *27*, 13273–13278.

Fernandez, J., Yaman, I., Huang, C., Liu, H., Lopez, A.B., Komar, A.A., Caprara, M.G., Merrick, W.C., Snider, M.D., Kaufman, R.J., et al. (2005). Ribosome stalling regulates IRES-mediated translation in eukaryotes, a parallel to prokaryotic attenuation. *Mol. Cell* *17*, 405–416.

Frey, U., and Morris, R.G. (1997). Synaptic tagging and long-term potentiation. *Nature* *385*, 533–536.

Giorgi, C., Yeo, G.W., Stone, M.E., Katz, D.B., Burge, C., Turrigiano, G., and Moore, M.J. (2007). The EJC factor eIF4AIII modulates synaptic strength and neuronal protein expression. *Cell* *130*, 179–191.

Gupta, K.C., and Ono, E. (1997). Stimulation of Sendai virus C' protein synthesis by cycloheximide. *Biochem. J.* *321*, 811–818.

Guzowski, J.F., McNaughton, B.L., Barnes, C.A., and Worley, P.F. (1999). Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles. *Nat. Neurosci.* *2*, 1120–1124.

Hou, L., Antion, M.D., Hu, D., Spencer, C.M., Paylor, R., and Klann, E. (2006). Dynamic translational and proteasomal regulation of fragile X mental retardation protein controls mGluR-dependent long-term depression. *Neuron* *51*, 441–454.

Huber, K.M., Kayser, M.S., and Bear, M.F. (2000). Role for rapid dendritic protein synthesis in hippocampal mGluR-dependent long-term depression. *Science* *288*, 1254–1257.

Huber, K.M., Gallagher, S.M., Warren, S.T., and Bear, M.F. (2002). Altered synaptic plasticity in a mouse model of fragile X mental retardation. *Proc. Natl. Acad. Sci. USA* *99*, 7746–7750.

Iacoangeli, A., Rozhdestvensky, T.S., Dolzhanskaya, N., Tournier, B., Schutt, J., Brosius, J., Denman, R.B., Khandjian, E.W., Kindler, S., and Tiedge, H. (2008). On BC1 RNA and the fragile X mental retardation protein. *Proc. Natl. Acad. Sci. USA* *105*, 734–739.

Ishizuka, A., Siomi, M.C., and Siomi, H. (2002). A Drosophila fragile X protein interacts with components of RNAi and ribosomal proteins. *Genes Dev.* *16*, 2497–2508.

Jin, P., Zarnescu, D.C., Ceman, S., Nakamoto, M., Mowrey, J., Jongens, T.A., Nelson, D.L., Moses, K., and Warren, S.T. (2004). Biochemical and genetic interaction between the fragile X mental retardation protein and the microRNA pathway. *Nat. Neurosci.* *7*, 113–117.

Kauderer, B.S., and Kandel, E.R. (2000). Capture of a protein synthesis-dependent component of long-term depression. *Proc. Natl. Acad. Sci. USA* *97*, 13342–13347.

Kelleher, R.J., 3rd, Govindarajan, A., and Tonegawa, S. (2004). Translational regulatory mechanisms in persistent forms of synaptic plasticity. *Neuron* *44*, 59–73.

Link, W., Konietzko, U., Kauselmann, G., Krug, M., Schwanke, B., Frey, U., and Kuhl, D. (1995). Somatodendritic expression of an immediate early gene is regulated by synaptic activity. *Proc. Natl. Acad. Sci. USA* *92*, 5734–5738.

Lyford, G.L., Yamagata, K., Kaufmann, W.E., Barnes, C.A., Sanders, L.K., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Lanahan, A.A., and Worley, P.F. (1995). Arc, a growth factor and activity-regulated gene, encodes a novel

- cytoskeleton-associated protein that is enriched in neuronal dendrites. *Neuron* 14, 433–445.
- Marrone, D.F., Schaner, M.J., McNaughton, B.L., Worley, P.F., and Barnes, C.A. (2008). Immediate-early gene expression at rest recapitulates recent experience. *J. Neurosci.* 28, 1030–1033.
- McBride, S.M., Choi, C.H., Wang, Y., Liebelt, D., Braunstein, E., Ferreira, D., Sehgal, A., Siwicki, K.K., Dockendorff, T.C., Nguyen, H.T., et al. (2005). Pharmacological rescue of synaptic plasticity, courtship behavior, and mushroom body defects in a *Drosophila* model of fragile X syndrome. *Neuron* 45, 753–764.
- Mori, Y., Imaizumi, K., Katayama, T., Yoneda, T., and Tohyama, M. (2000). Two cis-acting elements in the 3' untranslated region of alpha-CaMKII regulate its dendritic targeting. *Nat. Neurosci.* 3, 1079–1084.
- Muddashetty, R.S., Kelic, S., Gross, C., Xu, M., and Bassell, G.J. (2007). Dysregulated metabotropic glutamate receptor-dependent translation of AMPA receptor and postsynaptic density-95 mRNAs at synapses in a mouse model of fragile X syndrome. *J. Neurosci.* 27, 5338–5348.
- Nairn, A.C., and Palfrey, H.C. (1987). Identification of the major Mr 100,000 substrate for calmodulin-dependent protein kinase III in mammalian cells as elongation factor-2. *J. Biol. Chem.* 262, 17299–17303.
- Narayanan, U., Nalavadi, V., Nakamoto, M., Pallas, D.C., Ceman, S., Bassell, G.J., and Warren, S.T. (2007). FMRP phosphorylation reveals an immediate-early signaling pathway triggered by group I mGluR and mediated by PP2A. *J. Neurosci.* 27, 14349–14357.
- Nosyreva, E.D., and Huber, K.M. (2006). Metabotropic receptor-dependent long-term depression persists in the absence of protein synthesis in the mouse model of fragile X syndrome. *J. Neurophysiol.* 95, 3291–3295.
- Obrig, T.G., Culp, W.J., McKeenan, W.L., and Hardesty, B. (1971). The mechanism by which cycloheximide and related glutarimide antibiotics inhibit peptide synthesis on reticulocyte ribosomes. *J. Biol. Chem.* 246, 174–181.
- O'Donnell, W.T., and Warren, S.T. (2002). A decade of molecular studies of fragile X syndrome. *Annu. Rev. Neurosci.* 25, 315–338.
- Perlman, J., and Feldman, J.F. (1982). Cycloheximide and heat shock induce new polypeptide synthesis in *Neurospora crassa*. *Mol. Cell. Biol.* 2, 1167–1173.
- Pestova, T.V., Kolupaeva, V.G., Lomakin, I.B., Pilipenko, E.V., Shatsky, I.N., Agol, V.I., and Hellen, C.U. (2001). Molecular mechanisms of translation initiation in eukaryotes. *Proc. Natl. Acad. Sci. USA* 98, 7029–7036.
- Pinkstaff, J.K., Chappell, S.A., Mauro, V.P., Edelman, G.M., and Krushel, L.A. (2001). Internal initiation of translation of five dendritically localized neuronal mRNAs. *Proc. Natl. Acad. Sci. USA* 98, 2770–2775.
- Plath, N., Ohana, O., Dammermann, B., Errington, M.L., Schmitz, D., Gross, C., Mao, X., Engelsberg, A., Mahlke, C., Welzl, H., et al. (2006). Arc/Arg3.1 is essential for the consolidation of synaptic plasticity and memories. *Neuron* 52, 437–444.
- Ramirez-Amaya, V., Vazdarjanova, A., Mikhael, D., Rosi, S., Worley, P.F., and Barnes, C.A. (2005). Spatial exploration-induced Arc mRNA and protein expression: evidence for selective, network-specific reactivation. *J. Neurosci.* 25, 1761–1768.
- Rao, V.R., Pintchovski, S.A., Chin, J., Peebles, C.L., Mitra, S., and Finkbeiner, S. (2006). AMPA receptors regulate transcription of the plasticity-related immediate-early gene Arc. *Nat. Neurosci.* 9, 887–895.
- Ryazanov, A.G. (1987). Ca<sup>2+</sup>/calmodulin-dependent phosphorylation of elongation factor 2. *FEBS Lett.* 214, 331–334.
- Ryazanov, A.G. (2002). Elongation factor-2 kinase and its newly discovered relatives. *FEBS Lett.* 514, 26–29.
- Ryazanov, A.G., Shestakova, E.A., and Natapov, P.G. (1988). Phosphorylation of elongation factor 2 by EF-2 kinase affects rate of translation. *Nature* 334, 170–173.
- Schaeffer, C., Bardoni, B., Mandel, J.L., Ehresmann, B., Ehresmann, C., and Moine, H. (2001). The fragile X mental retardation protein binds specifically to its mRNA via a purine quartet motif. *EMBO J.* 20, 4803–4813.
- Scheetz, A.J., Nairn, A.C., and Constantine-Paton, M. (2000). NMDA receptor-mediated control of protein synthesis at developing synapses. *Nat. Neurosci.* 3, 211–216.
- Shepherd, J.D., Rumbaugh, G., Wu, J., Chowdhury, S., Plath, N., Kuhl, D., Huganir, R.L., and Worley, P.F. (2006). Arc/Arg3.1 mediates homeostatic synaptic scaling of AMPA receptors. *Neuron* 52, 475–484.
- Snyder, E.M., Philpot, B.D., Huber, K.M., Dong, X., Fallon, J.R., and Bear, M.F. (2001). Internalization of ionotropic glutamate receptors in response to mGluR activation. *Nat. Neurosci.* 4, 1079–1085.
- Steward, O., Wallace, C.S., Lyford, G.L., and Worley, P.F. (1998). Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites. *Neuron* 21, 741–751.
- Taubenfeld, S.M., Stevens, K.A., Pollonini, G., Ruggiero, J., and Alberini, C.M. (2002). Profound molecular changes following hippocampal slice preparation: loss of AMPA receptor subunits and uncoupled mRNA/protein expression. *J. Neurochem.* 81, 1348–1360.
- Tu, J.C., Xiao, B., Yuan, J.P., Lanahan, A.A., Leoffert, K., Li, M., Linden, D.J., and Worley, P.F. (1998). Homer binds a novel proline-rich motif and links group 1 metabotropic glutamate receptors with IP<sub>3</sub> receptors. *Neuron* 21, 717–726.
- Walden, W.E., and Thach, R.E. (1986). Translational control of gene expression in a normal fibroblast. Characterization of a subclass of mRNAs with unusual kinetic properties. *Biochemistry* 25, 2033–2041.
- Wang, H., and Tiedge, H. (2004). Translational control at the synapse. *Neuroscientist* 10, 456–466.
- Weiler, I.J., Spangler, C.C., Klintsova, A.Y., Grossman, A.W., Kim, S.H., Bertaina-Anglade, V., Khaliq, H., de Vries, F.E., Lambers, F.A., Hatia, F., et al. (2004). Fragile X mental retardation protein is necessary for neurotransmitter-activated protein translation at synapses. *Proc. Natl. Acad. Sci. USA* 101, 17504–17509.
- Westmark, C.J., and Malter, J.S. (2007). FMRP mediates mGluR5-dependent translation of amyloid precursor protein. *PLoS Biol.* 5, e52.
- Yuan, J.P., Kiselyov, K., Shin, D.M., Chen, J., Shcheynikov, N., Kang, S.H., Dehoff, M.H., Schwarz, M.K., Seeburg, P.H., Muallem, S., and Worley, P.F. (2003). Homer binds TRPC family channels and is required for gating of TRPC1 by IP<sub>3</sub> receptors. *Cell* 114, 777–789.
- Zalfa, F., Giorgi, M., Primerano, B., Moro, A., Di Penta, A., Reis, S., Oostra, B., and Bagni, C. (2003). The fragile X syndrome protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at synapses. *Cell* 112, 317–327.